

Biogeography of the Sacoglossa (Mollusca, Opisthobranchia)*

Kathe R. JENSEN¹⁾

¹⁾Zoological Museum, Copenhagen, Denmark

*Paper presented to the 2nd International Workshop on Opisthobranchia, ZFMK, Bonn, Germany, September 20th to 22nd, 2006

Abstract. The Sacoglossa (Mollusca, Opisthobranchia) comprise almost 400 nominal species level taxa. Of these 284 are considered valid (i.e., no published synonymies) in this study. About half of the nominal species have been described before 1950, and the 10 most productive taxonomists have described about half of the species. Distributions of all valid species are reviewed. The highest diversity is found in the islands of the Central Pacific, though species diversity is almost as high in the Indo-Malayan sub-province. The Caribbean forms another center of species diversity. These three areas are distinguished by the high number of Plakobranchoidea. Similarity among provinces is generally low. Endemism is high in most provinces, but this may be an artifact of collecting activity. The decrease in number of species with latitude is spectacular, and the number of cold-water endemics is very low, indicating that sacoglossans in cold temperate regions are mostly eurythermic warm water/ tropical species. The highest number of species in cold temperate areas is found in Japan and Southeastern Australia. This coincides with high species diversity of the algal genus *Caulerpa*, which constitutes the diet of all shelled and many non-shelled sacoglossans.

Keywords. Species diversity, endemism.

1. INTRODUCTION

Information on biogeography is important for understanding speciation and phylogeny as well as for making decisions about conservation. Ideally, combining a phylogenetic tree with a distributional map should give information on whether species dispersed from a center of origin or were the result of vicariance events. For most marine invertebrate groups, however, phylogenies are not fully resolved and/or taxonomy is not yet stable, and even information on distributions is incomplete. Species are still being split or synonymized, and new and undescribed species are discovered. In a worst case scenario a distribution map would show the activities of taxonomists rather than actual species distributions. In the present study existing distributional data for the Sacoglossa (Mollusca: Opisthobranchia) is reviewed and analyzed with regard to different biogeographic theories as well as activities of taxonomists over time. Phylogenetic analysis has been performed at the genus level (JENSEN 1996a), and for one genus, *Thuridilla*, at species level (GOSLINER 1995). The relationship of the Sacoglossa to other opisthobranchs has been discussed in several recent publications (JENSEN 1996b; MIKKELSEN 1996, 1998; THOLLESON 1999; WÄGELE et al. 2003).

Sacoglossans are suctorial herbivores; only two or three species are oophagous, feeding on the eggs of other opisthobranchs (JENSEN 1993a, 1997a). This means that

they have depth distributions restricted to the photic zone, i.e. generally <100m. Sacoglossans are also dietary specialists, the majority of species feeding on siphonaceous green algae, especially *Caulerpa* spp. (Jensen 1997a). Hence they only occur in the habitats where these algae are found. The total number of valid species is around 300, but new species are still described and other species are synonymized.

2. MATERIALS AND METHODS

Distributional data for all species of Sacoglossa were taken from the literature. The study has included most publications of original descriptions to get the type localities. However, in the case of the oldest descriptions, the publications by SCHMEKEL & PORTMANN (1982) and BOUCHET (1984) have been used. Also, national and regional faunal checklists have been included, as well as records published on the Sea Slug Forum (<http://www.seaslugforum.net/>). All nominal species listed in Appendix 1 have been included in the first analysis for bias of taxonomic expertise and scientific activity. In the distributional analyses, however, only species considered valid in this study have been included. As the present study is not a taxonomic analysis, species identifications and syn-

onymizations, with a few controversial exceptions, will not be discussed. Only synonymies that have been published and not subsequently contested are used. Thus species that have only been mentioned once in the literature are, with few exceptions mentioned in the text, considered valid.

Biogeographic regions and provinces were taken from BRIGGS (1995) (Fig. 1), and sacoglossan distributions among these provinces were recorded. Although it must be assumed that a species occurs continuously between the extreme points of distribution, species were only scored as occurring in a region or province if at least one published record existed. The number of endemic species was determined for each province. As some regions were clearly underrepresented with regards to faunistic studies on opisthobranchs, a few regions have been merged or deleted from the analyses. Similarity between biogeographic regions or provinces was analyzed using three indices: CJ= Jaccard's coefficient= $100(a/(N1+N2-a))$ (Valentine 1966), SD= Dice coefficient = $100(2a/(2a+b+c))$ (LEAL & BOUCHET 1991), and I=index

of inclusion= $100(a/Nmin)$ (GOLIKOV 1989). These indices differ in the weight placed on shared species (a) compared to total number of species in the compared regions (N1, N2), and species found exclusively in one or the other of the compared regions (b, c).

3. RESULTS

3.1. Fossil history

After the description of live specimens of bivalved sacoglossan gastropods (KAWAGUTI & BABA 1959), several papers on fossil species of these sacoglossans appeared. The first reviews of fossil sacoglossans were those of BOETTGER (1963) and KAY (1968). There have been extensive discussions about the identity of the Recent *Tamanovalva*, *Edentellina* and *Midorigai* and the Middle Eocene genus *Bertheliina* (e.g. EDMUNDS 1963; BURN 1998). KEEN & SMITH (1961) listed several other fossil species and included all in the family Juliidae Dall, 1898, which had previously been located in the Bivalvia. More




Fig. 1. Map showing biogeographic regions used in the present study. Regions have been modified from BRIGGS (1995). 1. North-east Atlantic. 2. Lusitanian. 3. Mediterranean (including the Black Sea). 4. Northwest Atlantic. 5. Caribbean. 6. Southwestern Atlantic tropical and warm temperate. 7. Aleutian. 8. Oregonian. 9. Californian. 10. Mexican-Panamanian. 11. Southeastern Pacific warm temperate. 12. Cold temperate South America. 13. Northwest Pacific cold temperate. 14. Northwest Pacific warm temperate. 15. Northern and northwestern Australia. 16. Southwestern and southern Australia. 17. Southeastern Australia. 18. Northeastern Australia. 19. Great Barrier Reef. 20. New Zealand. 21. Southeast Atlantic. 22. Western Indian Ocean. 23. Indo-Polynesian region (including Ryukyu Islands). 24. Hawaii.

recently several more fossil bivalved sacoglossans have been described (see LE RENARD et al. 1996 for review), and also a single species of *Volvatella* has been described from the Lower Miocene of France (VALDÉS & LOZOUET 2000). Thus there may be two or five Recent genera of bivalved sacoglossans, whereas there are 9 fossil genera extending from the Lower Eocene to Lower Pliocene. Most fossil species have been found in European localities, but a few are from the Caribbean, and one each from Australia and Indonesia. However, no doubt more fossil species will be described in the future.

The temporal and spatial distribution of fossil sacoglossans indicates that they arose as part of the Tethys Sea fauna. As sea level receded and temperatures cooled down, their distribution became more restricted, and today there is only one species of *Berthelinia* in the Caribbean and one in the Panamanian region; the remaining species are Indo-West Pacific. *Julia* has one species in the East Pacific; the remaining species are Indo-West Pacific. For *Volvatella* there is only one species in the Caribbean, one in warm temperate South Africa, and the remaining species are Indo-West Pacific. The disappearance of a major part of the coral reefs at the end of the Cretaceous (BRIGGS 1995) may have created ideal conditions for speciation of siphonaceous green algae when sea level rose again in early Eocene.

3.2. Recent species

Slightly more than half (199 of 387) of the nominal species have been described before 1950. There is a distinct peak around the 1860s and 1870s when PEASE and BERGH were most active describing species from the Indo-West Pacific and Costa and Trinchese worked in the Mediterranean (Fig. 2). After 1950, it is especially the MARCUSES (39 species) and K. BABA (30 species) who dominate the number of new species (Table 1). The 10 most productive authors or groups of authors have described almost 50 % of the species.

Of the 387 nominal species 284 (73%) have been included in the similarity analyses. The number of species recorded from regions and provinces shown in Fig. 1 is listed in Table 2. Some regions are distinctly underrepresented in regards to number of records. This is true for most of the southern cold temperate zone, but also for tropical East Atlantic and southern East Pacific. Most of the biogeographic regions and provinces are supported by the present study as indicated by the percentage of endemic species. The regions and provinces with less than 10% endemism will be discussed below.

The Northeast Atlantic and Mediterranean were the earliest studied areas. The number of species described dur-




Fig. 2. Frequency of species descriptions through time.

Table 1. Number of species described by different authors (called "group(s)", when with one or several co-authors) through time. All nominal taxa have been included.

Author	# species described
Marcus (& Marcus), 1955–1982	39
Baba (et al.), 1935–1959	30
Jensen (et al.), 1980–1999	23
Bergh, 1871–1905	20
Pease, 1860–1871	17
Ortea (et al.), 1981–2006	15
Trinchese, 1869–1895	14
Ichikawa, 1993	11
Thompson (et al.), 1973–1988	10
A. Costa, 1862–1876	10
Total for 10 authors (groups)	189
Total # nominal species	387
Percentage described by 10 most productive authors (groups)	49
Total # authors (groups)	104
Average # species per author (group)	3.7

ing the 19th century is high, but many species have subsequently been synonymized. New species are still discovered (ORTEA & TEMPLADO 1990; PERRONE 1990; CERVERA et al. 1991), old synonyms resurrected (CERVERA & LOPEZ-GONZÁLEZ 1996; ORTEA & MORO 1998), and the validity of some species, even some of the more recently described ones, is still debated (CERVERA et al. 2006). In addition, species ranges appear to be expanding (THOMPSON 1983; ORTEA et al. 1997; EVERTSEN & BAKKEN 2002). Hence the present analyses represent an *ad hoc* picture of species distributions and diversity.

Three faunal provinces are recognized in the Northeast Atlantic: the warm temperate Mediterranean Sea, including the Black and Azov Seas (no sacoglossans occur in the Caspian and Aral Seas), the warm temperate Lusitanian province and the cold temperate Northeast Atlantic Boreal region (BRIGGS 1995). Three species appear to be endemic to the Northeast Atlantic cold-water region. However, two of these may be identical to Lusitanian and/or Mediterranean species. The possible synonymy of *Ercolania nigra* and *E. viridis* is currently under study by the present author, and it is also likely that *Calliopaea oopha-*

Table 2. Species distribution and endemism of sacoglossan opisthobranch in biogeographic regions as defined by BRIGGS (1995). Only species considered valid in the present study have been included. n.d. not determined.

Region	# Species	# endemics (%)
1. Northeast Atlantic boreal	11	3* (27)
2. Lusitanian	40	5 (12.5)
3. Mediterranean + Black Sea	37	8 (22)
4. Northwest Atlantic boreal	6	1 (17)
5. Caribbean incl. Florida	49	21 (43)
6. Southwestern Atlantic tropical + warm temperate	19	3 (16)
7. Aleutian	6	0
8. Oregonian	8	0
9. Californian	9	0
10. Mexican-Panamanian	23	9 (39)
11. Southeast Pacific warm temperate	6	0
12. Cold temperate South America	4	2 (50)
13. Northwest Pacific cold temperate	29	5 (17)
14. Northwest Pacific warm temperate	41	9 (21)
15. North and Northwestern Australia	23	4 (17)
16. South and Southwestern Australia	22	6 (27)
17. Southeastern Australia	15	6 (40)
18. Northeastern Australia	18	1 (5.5)
19. Great Barrier Reef	22	0
20. New Zealand	5	1 (20)
21. Southeastern Atlantic	9	4 (44)
22. Western Indian Ocean + Red Sea	32	8 (25)
23. Indo-Polynesian, incl. Ryukyu Islands	107	n.d.
24. Hawaii	25	6 (24)

*Two of these species may be synonymous with Lusitanian and Mediterranean species.

Table 3. Similarity between provinces of the Atlantic Ocean. The Southeast Atlantic is excluded due to lack of information. In this and all following tables only species considered valid in the present study have been included. CJ Jaccard's Coefficient; SD Dice Coefficient; I Index of inclusion; N number of species included; n.d. not determined.

a.						
CJ \ SD	NE Atl cold N=11	NW Atl cold N=6	Lusitan. N=40	Medit N=38	Carib N=49	Brazil N=19
NE Atl cold	-	23.5	27.5	28.6	3.33	n.d.
NW Atl cold	13.3	-	8.70	9.09	14.5	n.d.
Lusitanian	15.9	4.55	-	66.7	31.5	13.6
Mediterranean	16.7	4.76	50.0	-	23.0	n.d.
Caribbean	1.69	7.84	18.7	13.0	-	47.1
Brazil	n.d.	n.d.	7.27	n.d.	30.8	-

b.						
I	NE Atl cold	NW Atl cold	Lusitanian	Medit	Carib	
NE Atl cold	-	-	-	-	-	
NW Atl cold	33.3	-	-	-	-	
Lusitanian	63.6	33.3	-	-	-	
Mediterranean	63.6	33.3	68.4	-	-	
Caribbean	9.09	66.7	35.0	26.3	-	
Brazil	n.d.	n.d.	21.1	n.d.	84.2	

ga Lemche, 1974 is a subtidal variety of *C. bellula* d'Orbigny, 1837. *Limapontia depressa* may be the only endemic from this cold-water region, and even that species may occasionally be found further south on the French Atlantic coast (M. PODDUBETSKAIA, Bordeaux, pers. comm. 2006). The other two species of *Limapontia* only extend into the Lusitanian and/or Mediterranean province (PRUVOT-FOL 1954; SCHMEKEL & PORTMANN 1982); *L. capitata* occurs in all three provinces and *L. senestra* occurs in two provinces. *L. capitata* is also the only saeoglossan recorded from Iceland and the Faroe Islands (PLATTS 1985). Hence the genus *Limapontia* is probably a cold-water, eurythermal genus, which is endemic to the Northeast Atlantic-Mediterranean region. Various records exist of unidentified species of *Limapontia* from other regions (e.g. ENGEL et al. 1940; BURN 1973; SCHRÖDL 1996). However, these need confirmation as many species of *Ercolania laevis* in juveniles.

One species, *Hermaea variopicta*, has its northern limit along the south coast of the UK (LEMICHE & THOMPSON 1974). Two species, *Elysia viridis* and *Calliopa bellula*, have their northern limit around Trondheimfjord (BRATTEGARD & HOLTHE 2001). Four species, *Alderia modesta*, *Placida dendritica*, *Limapontia capitata* and *L. senestra*, occur in the northernmost part of Norway (VADER 1981; BRATTEGARD & HOLTHE 2001; EVERTSEN & BAKKEN 2002; pers. obs.), and all three species of *Limapontia* plus *Alderia modesta* have been recorded from the White Sea (ROGINSKAYA 2000; MARTYNOV et al. 2006). Prior to 1997

P. dendritica had not been recorded from Norway north of the Bergen area (EVERTSEN & BAKKEN 2002).

Five species appear to be endemic to the Lusitanian province. These have all been described after 1980, so it is possible that they will be found in neighboring regions in the future. The Canary Islands has the highest species diversity of this region (27 of 40 species have been recorded here). A number of Caribbean species have been recorded from these islands in recent years (ORTEA et al. 1998). Since the saeoglossan fauna of these islands has been well documented over many years (ORTEA 1981; FERNANDEZ-OVIES & ORTEA 1986; CERVERA et al. 1988; ORTEA et al. 1990; TEMPLADO et al. 1990), and only one or a few specimens have been collected at one single time far from their native distribution area, these species have most likely been transported by human activities (CHAPMAN & CARLTON 1991). Few species have been recorded from Madeira, Salvage, Cape Verde and Azores Islands (ORTEA 1981; ORTEA & TEMPLADO 1990; ORTEA et al. 1988, 1990, 1998; MALAQUIAS & CALADO 1997; JENSEN 1995, in prep.). There is an old record of the shelled *Ascobulla fragilis* from the Atlantic coast of Spain, which is cited in more recent publications (PILSBRY 1895; PRUVOT-FOL 1954; CERVERA et al. 2006). As this species feeds exclusively on *Caulerpa*, which does not extend this far north (DOUMENGE 1995), this needs to be re-examined.

Eight species are endemic to the Mediterranean. Some of these may be synonyms of other species with wider dis-

tributions (SCHEMEKEL & PORTMANN 1982; THOMPSON 1988). Eight ampho-Atlantic (one of which may be cosmopolitan), eight Northeast Atlantic-Lusitanian, and five endemic species do not extend into the eastern basin of the Mediterranean. Only four species found in the eastern basin of the Mediterranean do not occur in the western basin; two of these have been described recently (THOMPSON 1988). The total number of sacoglossans in the eastern basin is only 17 whereas 34 species are known from the western basin (SWENNEN 1961; BARASH & DANIN 1971; SCHEMEKEL & PORTMANN 1982; THOMPSON 1983; BOUCHET 1984; THOMPSON et al. 1985; THOMPSON 1988; THOMPSON & JAKLIN 1988; CATTANEO-VIETTI & THOMPSON 1989; CERVERA et al. 2006). The southern coast of the Mediterranean has been insufficiently studied. It is uncertain whether two or three species extend into the Black and Azov Seas. *Limapontia capitata* and *Calliopaea bellula* (as *Stiliger bellulus*) have been recorded previously (MURINA & ARTEMIJEVA 1997), but recent pictures on the Sea Slug Forum (KURAKIN 2002) have shown that *Ercolania viridis* is present, and it remains to be seen whether the species previously identified as *C. bellula* has been misidentified, or whether both species occur.

The Lusitanian and Mediterranean provinces have very high similarity indices (Table 3). The combined number of species amounts to 52, which is very close to the species diversity found in the Caribbean. However, the number of Plakobranchidae is lower in the eastern Atlantic provinces (Table 9).

BRIGGS (1995) recognizes one circumpolar Arctic region comprising Spitzbergen, Greenland and the northern coasts of North America and Russia. No sacoglossans have been recorded from Spitzbergen (GULLIKSEN et al. 1999) or the north coast of North America (BLEAKNEY 1996; GODDARD & FOSTER 2002). The four species occurring in the White Sea also occur in the Russian part of the Barents Sea (ROGINSKAYA 2000; MARTYNOV et al. 2006). In addition, MARTYNOV et al. (2006) mentions an old record of a single juvenile specimen of *Placida dendritica* from Kola Bay. This indicates that this area is influenced by the North Atlantic Current and should be included in the boreal region. The single specimen of *Alderia modesta* recorded from western Greenland (PLATTS 1985) could be attributed to larvae transported from Canada, which may occasionally be able to find suitable habitats for metamorphosis in Greenland. The latitude of Disko Fjord is about the same as northern Norway, where the species occurs regularly. It should be mentioned that in Danish waters this species seems to have disappeared from localities where prior to 1997 it was abundant (pers. obs.). Whether this is due to habitat deterioration or increased temperature is unknown.

The fauna of the Northwest Atlantic is also very well studied (e.g. MARCUS & MARCUS 1970; MARCUS 1972a,b; MARCUS & HUGHES 1974; CLARK 1975; JENSEN & CLARK 1983; BLEAKNEY 1996), though new species are still being described from the tropical waters (ORTEA & ESPINOSA 1996, 2000, 2001, 2002; CABALLER et al. 2006; PIERCE et al. 2006). Only 6 species occur in the cold temperate province; one of these, *Placida dendritica*, may be cosmopolitan (BLEAKNEY 1989) and one, *Alderia modesta*, is circum-boreal. Apparently only one species, *Elysia catulus*, is endemic to the Northwest Atlantic cold water region (CLARK 1975). This species feeds on the seagrass *Zostera marina*, which does not occur in Florida. It is possible that *E. catulus* is a dark pigmented variety of the other seagrass feeding species, *Elysia serca* (JENSEN 1982), in which case there will be no endemic species for the Northwest Atlantic. *Ercolania fuscata* may occur from Nova Scotia, Canada to Sao Paulo, Brazil, but this distribution is based on synonymization with *E. vauellus* and *E. talis* (JENSEN & CLARK 1983). Two species, *Elysia chlorotica* and *Hermatea cruciata*, have their southern limit in Florida (JENSEN & CLARK 1983), i.e., just south of the cold-water region. The former species also occurs in the northern part of the Gulf of Mexico (BOONE 1982), and the latter has its northern limit in Massachusetts (MARCUS 1972a). The species presently known as *Limapontia zonata*, and known only from its original collection (GOULD & BINNEY 1870), is probably a flatworm; no sacoglossan has transverse pigment bands.

A separate Carolinian province could not be distinguished for sacoglossan opisthobranchs, and no difference is evident between the continental and insular parts of the Caribbean (CLARK & DEFRESE 1987), including Bermuda (CLARK 1984), although these provinces were considered distinct by BRIGGS (1995). However, the majority of studies involving sacoglossans are from the Caribbean islands. The limit between cold-water and tropical faunas appears to be along the coast of Florida (JENSEN & CLARK 1983); 36 of the 49 Caribbean species have been recorded from Florida (THOMPSON 1977; MARCUS 1977, 1980; CLARK 1982; CLARK & DEFRESE 1987; PIERCE et al. 2006; VALDÉS et al. 2006). The degree of endemism in the Caribbean is exceptionally high (Table 2). Especially the number of Plakobranchidae is high (Table 9), indicating that speciation in this family has taken place within the province. Five of the 21 endemic species have been described after 1990, so they may be found outside this province in the future or be synonymized with other species. As for the Mediterranean and Lusitanian provinces, synonymies are extensively debated and the status of several species remains uncertain (JENSEN & CLARK 1983; MARCUS 1980; VALDÉS et al. 2006).

Table 4. Similarity of provinces of East Pacific. Due to the high similarity between Aleutian, Oregonian and Californian provinces, these have been merged (Al+Or+Cal). Other abbreviations as in Table 3.

a.						
CJ \ SD	Aleutian N=6	Oregonian N=8	Californian N=9	Al+Or+Cal N=10	Mex-Panam N=23	SE Pacific N=9
Aleutian	-	85.7	66.7	-	27.6	n.d.
Oregonian	75.0	-	82.4	-	38.7	n.d.
Californian	50.0	70.0	-	-	50.0	n.d.
Al+Or+Cal	-	-	-	-	48.5	10.5
Mex-Panam	16.0	24.0	33.3	32.0	-	43.8
SE Pacific	n.d.	n.d.	n.d.	5.56	28.0	-

b.					
I	Aleutian	Oregonian	Californian	Al+Or+Cal	Mex-Panam
Aleutian	-	-	-	-	-
Oregonian	100.0	-	-	-	-
Californian	83.3	87.5	-	-	-
Mex-Panam	66.7	75.0	88.9	80.0	-
SE Pacific	n.d.	n.d.	n.d.	11.1	77.8

The genus *Bosellia* appears to be an Atlantic warm-water genus with one ampho-Atlantic, one Canary Islands endemic (FERNANDEZ-OVIES & ORTEA 1986), and two Caribbean species (MARCUS 1973). There are some scattered reports of *Bosellia* from the Indo-Pacific region (MARCUS 1978; IMAMOTO 2004; PITTMAN 2004; RIEK 2006). However, too few specimens have been recorded to either identify them as one of the described species or decide that they are undescribed species.

BRIGGS (1995) recognized a tropical Brazilian province extending to just south of Rio de Janeiro. For sacoglossans, most Brazilian species extend south to the area around Sao Paulo (MARCUS (ER.) 1955, 1957; MARCUS & MARCUS 1967; MARCUS (EV.) 1977). This can probably be explained by the extensive collecting activity of the Marcuses around Sao Paulo; 47% of the Brazilian species have been described by them. Furthermore, more than 90% of the Brazilian species also occur in the Caribbean (Table 3b), and the number of endemic species is low (Table 2). This is most likely also due to the activities of the Marcuses in both these regions.

TROWBRIDGE (2002) reviewed the Northeast Pacific sacoglossan fauna. She recognized four provinces, but unfortunately the borders are not exactly the same as suggested by BRIGGS (1995). This is especially evident for the Californian province, in which Trowbridge records one species of the bivalved genus *Berthelinia* plus a couple of unidentified/undescribed species. The present study found no endemic species in the Aleutian, Oregonian and Californian provinces and hence these were merged. Also, no difference was obvious between the Mexican and Panamanian provinces, which have also been merged before

comparisons with other provinces (Table 4). In the Mexican-Panamanian province four species have been recorded after the publication of Trowbridge's paper (BEHRENS & HERMOSILLO 2005; KRUG et al. 2007) and one species, *Ascobulla californica* (originally described as *Cylindrobulla californica* by HAMATANI (1971)), was not considered a sacoglossan by TROWBRIDGE (2002). The occurrence of *Alderia modesta* in the Mexican-Panamanian region is probably the recently described species, *Alderia willowi* (KRUG et al. 2007), which occurs southwards from central California. The monotypic genus *Olea* is endemic to the Northeastern Pacific region, extending from the Aleutian to the Californian province (TROWBRIDGE 2002). *Hermatea vancouverensis* is a cold-water species, extending across the Bering Strait to the Kurile province (CHERNYSHEV 2005). One species, *Elysia hedgpethi*, occurs from British Columbia, Canada to Chile (SCHRÖDL 1996; TROWBRIDGE 2002), though the occurrence in Chile needs verification.

The species extending into the warm temperate region of the Southeast Pacific almost all are shared with the tropical Mexican-Panamanian region (SCHRÖDL 1996; TROWBRIDGE 2002; BEHRENS & HERMOSILLO 2005). *Julia thecaphora* is considered the oldest name for *J. equatorialis*, which was also described from the tropical East Pacific. Only four species have been recorded from cold-temperate South America, one from the Atlantic coast and three from the Pacific (MARCUS 1959; SCHRÖDL 1996; MUNIAN & ORTEA 1997). Due to the low number of species and sparse collecting activity, the species from the Southeast Pacific coast have been considered as one province for analyses.

Table 5. Similarity of Japanese biogeographic provinces and of Northwest and Northeast Pacific provinces. The Japanese provinces have also been compared to the neighboring Central Pacific sub-province. Abbreviations: see Table 3.

a.						
CJ \ SD	Japan cold temp., N=26	Japan warm temp., N=41	Ryukyu N=33	Central Pacif. N=51	NW Pacif. cold temp., N=29	NE Pacif. cold temp. N=10
Japanese cold temp.	-	59.7	27.1	26.0	-	n.d.
Japanese warm temp.	42.6	-	37.8	37.0	n.d.	n.d.
Ryukyu	15.7	23.3	-	38.1	n.d.	n.d.
Central Pacific	14.9	22.7	23.5	-	n.d.	n.d.
NW Pacific cold temp.	-	n.d.	n.d.	n.d.	-	20.5
NE Pacific cold temp.	n.d.	n.d.	n.d.	n.d.	11.4	-
b.						
I	Japan cold temp.	Japan warm temp.	Ryukyu	NW Pacif. cold temp.		
Japanese cold temp.	-	-	-	-		
Japanese warm temp.	76.9	-	-	-		
Ryukyu	30.8	42.4	-	-		
Central Pacific	38.5	41.5	48.5	-		
NE Pacific cold temp.	-	n.d.	n.d.	40.0		

Four biogeographic regions or provinces can be distinguished along the coasts of Japan. Biogeographically the southernmost archipelago of Ryukyu belongs in the vast tropical Indo-Polynesian province (BRIGGS 1995), but being under Japanese jurisdiction, the saeoglossan fauna has been studied mostly by Japanese scientists (e.g. BABA 1936; HAMATANI 1980; ICHIKAWA 1993). Hence Ryukyu has been included in both analyses of the Japanese fauna (Table 5) and of the Indo-West Pacific one (Table 7). The warm-water temperate region comprises southern and eastern Japan, including the well studied Seto Inland Sea and Sagami Bay. The corresponding continental coast of China has been insufficiently studied, and only 14 species have been recorded from southern Korea (KOH 2002a,b, 2003, 2005a,b,e; RUDMAN, Sydney, pers. comm. 2007). The cold-water temperate oriental province includes the central and western coasts of Honshu, whereas the northernmost island of Hokkaido belongs to the Kurile province (BRIGGS 1995). The cold-temperate fauna of Japan contains more species (N=26) than any other cold-water fauna. A few shelled species extend into this province, which is also seen in southern Australia, but not in other cold-water provinces. Only three additional saeoglossan species have been recorded from the Kurile province (BABA 1935; CHERNYSHEV & CHABAN 2005; TROWBRIDGE 2006) and these were only included in the comparison of Northeast and Northwest Pacific cold-water faunas (Table 5). The highest number of species has been recorded from the warm temperate region (e.g. BABA 1949, 1952a,b, 1955, 1957, 1959, 1966; 1968; HAMATANI 1968, 1969, 1972, 1976a,b, 1994; KAWAGUTI

& BABA 1959; HIRANO et al. 2006); in fact BABA described 59% of the species from this province and 69% of the species from the cold-water region of Japan. A few of the species described by BABA have been synonymized with more widespread Indo-West Pacific species (BABA 1974; JENSEN 1985). On the other hand, many of BABA's species have been identified outside Japan (JENSEN 1985; CARLSON & HOFF 1978, 2003; BURN 1998, 2006).

The Northeast Australian and Great Barrier Reef faunas (BURN 1966b; THOMPSON 1973; MARSHALL & WILLAN 1999; WÄGELE & JOHNSON 2001) are so closely related to each other that no endemics have been recorded from the Great Barrier Reef and only one endemic species, *Placida fralila*, has been recorded from Northeast Australia (Table 2). Hence these two provinces were merged for similarity analyses, and the combined province has two endemic species (*Elysia bennettiae*, *P. fralila*). For the Great Barrier Reef 30% of the species were listed as unidentified and/or undescribed (MARSHALL & WILLAN 1999). This part of Australia has been included in the Indo-Polynesian province by BRIGGS (1995). This is supported by the low endemicity, and also, the similarity with the fauna of the South Pacific islands is higher than with any of the other Australian provinces (Tables 6 and 7). The North and Northwestern Australian fauna has rather high similarity to the Western Indian Ocean fauna (Table 7). This is in spite of the fact that almost 40% of the species have been described by the present author within the last 20 years (JENSEN 1993b, 1997b,c; JENSEN & WELLS 1990). The fauna of the South and Southwestern Australia has

Table 6. Similarity of provinces of the Australian continent. Due to the high similarity between the Northeast Australian province and the Great Barrier Reef, these two provinces have been merged (NE Aus+GBR). Abbreviations as in Tab. 3.

a.						
CJ \ SD	NE Austr N=18	GBR N=22	NE Aus+ GBR N=28	N+NW Austr N=23	SW+S Austr N=22	SE Austr N=15
NE Australia	-	60.0	-	34.1	30.0	30.3
GBR	42.9	-	-	26.7	18.2	16.2
NE Aus+GBR	-	-	-	31.4	28.0	27.9
N+NW Austr	20.6	15.4	19.0	-	48.8	26.3
SW+S Austr	17.6	10.0	16.3	32.4	-	43.2
SE Austr	17.9	8.82	16.2	15.2	27.6	-

b.					
I	NE Austr	GBR	NE Aus+GBR	N+NW Austr	SW+S Austr
NE Australia	-	-	-	-	-
GBR	66.7	-	-	-	-
NE Aus+GBR	-	-	-	-	-
N+NW Austr	38.9	27.3	34.8	-	-
SW+S Austr	33.3	18.2	31.8	50.0	-
SE Austr	33.3	20.0	40.0	33.3	53.3

the highest affinity to the fauna of North and Northwestern Australia, and the other way around, whereas the fauna of Southeastern Australia (BURN 1958, 1960, 1965, 1974, 1998, 2006) has a higher affinity to the fauna of South and Southwestern Australia than to that of Northeastern Australia and the Great Barrier Reef (Table 6). This may change when the 50% unidentified/undescribed species listed for Southeastern Australia (BURN 2006) are properly named.

Very few sacoglossans have been recorded from New Zealand (POWELL 1937; WILLAN & MORTON 1984; TROWBRIDGE 1995a; SPENCER & WILLAN 1995). The fauna consists of widespread species and one endemic (Table 2). Hence this fauna has not been further analyzed in the present study.

The sacoglossan fauna of the Red Sea (ELIOT 1908; O'DONOGHUE 1929; HELLER & THOMPSON 1983) has about the same affinity to the fauna of the Indian subcontinent as to the Western Indian Ocean, and the index of inclusiveness (I) for the Red Sea and India *sensu lato* (*s.l.*, see below) is twice that for the Red Sea and the Western Indian Ocean (Table 7b). Also, two out of the ten recorded species are endemic to the Red Sea. Hence the Red Sea should be considered a separate province. In other groups of invertebrates Lessepsian migrants are common. For sacoglossans this appears to be a small and recent problem (YOKES 2001, 2002; RUDMAN 2002). Except for these few species, the Red Sea does not share any species with the Mediterranean.

The sacoglossans found in southwestern Africa are more closely related to those found in southeastern Africa than to any other region or province (GOSLINER 1987a); in fact no species are shared with the Brazilian fauna and only one species, *Placida dendritica*, is shared with the Lusitanian province. Hence this province has been considered in connection with the Western Indian Ocean province. Endemicity is high, but this could be due to the poor knowledge of tropical western Africa. Only one species has been described from this region (MARCUS & MARCUS 1966), so this was not included in the present study. The East Atlantic species *Elysia viridis* apparently occurs from central Norway (BRATTEGARD & HOLTHE 2001) to South Africa (GOSLINER 1987a), though no records exist between the Senegal (PRUVOT-FOL 1953) and South Africa, and GOSLINER (1998) has subsequently changed the identification to *Elysia* sp. The species was originally identified in South Africa as the Indian species *E. punctata* by MACNAE (1954). GOSLINER (1987a) found a distinct faunal separation for opisthobranchs at Port Elizabeth, whereas BRIGGS (1995) considers the coast between the Cape of Good Hope and north of Durban one province. In the present study Port Elizabeth has been used to separate the faunas of southwestern Africa and the Western Indian Ocean.

The Western Indian Ocean is considered a separate province by BRIGGS (1995). The sacoglossan fauna of this province has a high similarity with the South Pacific and North and Northwestern Australia (Table 7). The affinity with the fauna of India *s.l.* (India, Sri Lanka and Maldives) is considerably lower, and also the affinity with the Indo-

Table 7. Similarity of provinces of the Australian continent. Due to the high similarity between the Northeast Australian province and the Great Barrier Reef, these two provinces have been merged (NE Aus+GBR). Abbreviations as in Tab. 3.

a.

CJ \ SD	SW Afr. N=8	WIO N=27	Red Sea N=10	India s. I. N=20	Indo-Malay N=48	N+NW Austr N=23	Centr. Pac. isl. N=51	NE Aus + GBR N=28	S Pacific N=33	Hawaii N=25	Ryukyu N=33
SW Afr.	-	28.6	11.1	7.14	n.d.	12.9	n.d.	n.d.	9.76	n.d.	n.d.
WIO	16.7	-	27.0	17.0	24.0	32.0	35.9	25.5	36.7	n.d.	n.d.
Red Sea	5.88	15.6	-	26.7	n.d.	18.2	n.d.	n.d.	18.6	n.d.	n.d.
India s. I.	3.70	9.30	15.4	- 20.6	14.0	19.7	12.5	22.6	17.8	7.55	
Indo-Malay	n.d.	13.6	n.d.	11.5	-	22.5	50.5	28.9	37.0	19.2	24.7
N+NW Austr	6.90	19.0	10.0	7.50	12.7	-	29.7	31.4	32.1	n.d.	n.d.
Centr. Pac. isl.	n.d.	21.9	n.d.	10.9	33.8	17.5	-	43.0	50.0	42.1	38.1
NE Aus+GBR	n.d.	14.6	n.d.	6.67	16.9	19.0	27.4	-	42.6	n.d.	n.d.
S Pacific	5.13	22.4	10.3	12.8	22.7	19.1	33.3	27.1	-	41.2	24.2
Hawaii	n.d.	n.d.	n.d.	9.76	10.6	n.d.	26.7	n.d.	26.1	-	13.8
Ryukyu	n.d.	n.d.	n.d.	3.92	14.1	n.d.	23.5	n.d.	13.8	7.41	-

b.

I	SW Afr.	WIO	Red Sea	India s. I.	Indo-Malay	N+NW Austr.	Centr. Pac. isl.	NE Aus + GBR	S Pacific	Hawaii
SW Afr.	-	-	-	- -	-	-	-	-	-	-
WIO	62.5	-	-	- -	-	-	-	-	-	-
Red Sea	12.5	50.0	-	- -	-	-	-	-	-	-
India s. I.	12.5	20.0	40.0	- -	-	-	-	-	-	-
Indo-Malay	n.d.	33.3	n.d.	35.0	-	-	-	-	-	-
N+NW Austr	25.0	34.9	30.0	15.0	34.8	-	-	-	-	-
Centr. Pac. isl.	n.d.	51.9	n.d.	35.0	52.1	47.8	-	-	-	-
NE Aus+GBR	n.d.	25.9	n.d.	15.0	39.3	34.8	60.7	-	-	-
S Pacific	25.0	40.7	40.0	30.0	45.5	39.1	63.6	46.4	-	-
Hawaii	n.d.	n.d.	n.d.	20.0	28.0	n.d.	64.0	n.d.	48.0	-
Ryukyu	n.d.	n.d.	n.d.	10.0	30.3	n.d.	48.5	n.d.	24.2	16.0

Malayan sub-province is lower. Endemicity is high (5 of 27 species), though two of these species have sometimes been synonymized with widespread Indo-West Pacific species (GOSLINER 1987b). The majority of species recorded from the Western Indian Ocean have been collected from the southern part of the region, i.e. South Africa (THOMPSON 1979; GOSLINER 1987a,b, 1995), Tanzania (ELIOT 1903, 1904; GOSLINER 1995), Madagascar (GOSLINER 1995) and Mauritius (BERGH 1888; GOSLINER 1995).

A total of 107 species of sacoglossans have been recorded from the vast Indo-Polynesian province. Most of these species are only distributed in part of the province and hence it was subdivided into five sub-provinces: The Indian subcontinent, including Sri Lanka and the Maldiv Islands was considered one sub-province (10 endemic species); the Andaman Sea, the South China Sea, Indonesia and the Philippines form an Indo-Malayan sub-

province (9 endemic species); the Mariana and Marshall Islands together with Micronesia were considered a Central Pacific sub-province (4 endemic species); Papua New Guinea, Solomon Islands, Fiji, Vanuatu, New Caledonia, Samoa and the Polynesian islands form a South Pacific province (6 endemic species); and as mentioned above the Ryukyu Islands form a separate sub-province (10 endemic species). Many of the endemic species have been described within the last 20 years, so they may actually have wider distributions.

Of the 107 species recorded from the Indo-Polynesian province only 12 have distributions from the Western Indian Ocean and/or the Red Sea to the Central and/or South Pacific islands, two species, *Elysia ornata* and *Ercolania coerulea*, are circum-tropical and one, *Placida dendritica*, may be cosmopolitan. Two of the 12 widespread Indo-West Pacific species are shelled (*Oxynoe viridis* and *Berthelinia schlumbergeri*), seven are plakobranchoids (5

species of *Thuridilla*, *Elysiella psilla* and *Plakobranchms ocellatus*) and three are limapontioids (*Cyerce elegans*, *C. nigricans* and *Polybranchia orientalis*). The widest latitudinal distributions along the West Pacific rim are found in the four species that occur from the cold-temperate part of Japan to the cold-temperate part of Australia. Two of these species are the circum-tropical *Elysia ornata* and the questionably cosmopolitan *Placida dendritica*; the other two are *Elysia obtusa* and *Stiliger smaragdinus*. *P. dendritica* has been synonymized with a number of species described from different places in the Indo-West Pacific (BLEAKNEY 1989), but the synonymy has been doubted (e.g. TROWBRIDGE 1995b). Two species, *Oxyroe viridis* and *Elysiella psilla*, extend from warm-temperate Japan to southern Australia. Two further species, *Thuridilla splendens* and *Polybranchia orientalis*, extend from warm-temperate Japan to Northeast Australia and the Great Barrier Reef. Four species, *Plakobranchms ocellatus*, *Thuridilla vatae*, *T. hoffae* and *Cyerce nigricans*, extend from the Ryukyu Islands to tropical Australia. Most of these species also have wide longitudinal distributions. *Plakobranchms ocellatus* occurs in a number of colour varieties, and it is possible that a complex of sibling species is involved (see discussion on the Sea Slug Forum: <http://www.seaslugforum.net/find.cfm?id=13970>) (last access 12th of August 2007).

The Indian sub-province has a high proportion of endemic species (50%). This is probably an artifact due to the activities of a few taxonomists who have worked only within this sub-province (KELAART 1858; NEVILL & NEVILL 1869; RAO 1937; RAO & RAO 1963; SARMA 1975). Most of G. & H. NEVILL's species have been synonymized, though not consistently with the same species. Although doubtful, they have been considered valid in the present study. The species described by KELAART have been re-examined several times (ELIOT 1906; O'DONOGHUE 1932) and they are still recognized as valid, mostly widespread Indo-West Pacific species. In spite of the high endemicity, *India s.l.* shows higher similarity to the Indo-Malayan, Central and South Pacific sub-provinces than to the Western Indian Ocean province (Table 7). The highest similarity is found between *India s.l.* and the Red Sea, but this is caused by the low number of species found in the Red Sea, and the high proportion of widespread Indo-West Pacific species.

The Indo-Malayan and Central Pacific sub-provinces have twice the number of species occurring in tropical Australia and the Western Indian Ocean and 50% more species than the South Pacific and Ryukyu Islands. This could be seen as evidence for being a center of origin for evolution of new species. Looking at species composition in the central Pacific islands (data from MARCUS 1965; JOHNSON & BOUCHER 1983; CARLSON & HOFF 2003), it seems more

likely that they are "traps", where species dispersing from the Japanese warm-temperate and from tropical and possibly even warm-temperate Australia can find suitable habitats. The Central Pacific sub-province is the only one with less than 10% endemics, but many unidentified and/or undescribed species are known from this sub-province (CARLSON & HOFF 2003; own obs.). Most biogeographic studies indicate that the triangle consisting of Indonesia, Malaysia and the Philippines and sometimes including Papua New Guinea has the highest number of species (EKMAN 1953; BRIGGS 1995, 2005). Information on sacoglossan distributions in the Indo-Malayan sub-province has been collected from numerous sources (e.g. BERGH 1871, 1872, 1905; ELIOT 1917; LIN 1986, 1990; GOSLINER 1995; GOSLINER et al. 1996; DEBELIUS 1996; JENSEN 1998a,b, 2003; SWENNEN et al. 2001). The number of sacoglossan species is slightly lower in this sub-province than in the islands of the Central West Pacific, i.e. Marianas, Marshall Islands and Micronesia. This may be a collecting artifact caused by the activities of Carlson and Hoff in Guam and neighboring islands; they recorded 91 species of which 48 (53%) were identified to species level (CARLSON & HOFF 2003). However, the present similarity analyses also indicate that these islands constitute the center of species diversity. Species described from northern Australia, the South China Sea and southern Japan also occur in the Mariana Islands. The genus *Gascoignella* is endemic to the Indo-Malayan province (JENSEN 1985; SWENNEN 2001), and the genus *Sohgenia* is endemic to the Central Pacific islands (HAMATANI 1991). However, with one exception, these species have been described within the last 20 years, so they may have wider distributions.

BRIGGS (1995) considered the Hawaiian islands a separate region. The Hawaiian islands have a high endemicity (OSTERGAARD 1955; KAY 1967; present study), but the present study has shown that they also have a rather high similarity to the other islands of the Central and South Pacific (Table 7).

The islands of the South Pacific have been rather sporadically studied (PEASE 1861, 1866, 1868, 1871; ELIOT 1899; RISBEC 1928, 1953; BURN 1966a; MILLER 1969; BRODIE & BRODIE 1990; GOSLINER 1995). Hence the total number of species as well as the number of endemics may be considerably higher. GOSLINER & DRAHEIM (1996) estimated that more than 40% of the opisthobranch species from Papua New Guinea are undescribed. For the Fiji Islands 30% of the species have not been identified to species level (BRODIE & BRODIE 1990). For the sacoglossans most of the old species are poorly described and need re-examination.

4. DISCUSSION

Sacoglossans are small, often cryptically colored species and they are therefore difficult to collect. This means that only a few biogeographic provinces have been thoroughly studied. In recent years international workshops and biodiversity programs have greatly increased the number of sacoglossans known from more remote localities, especially in tropical waters (JENSEN 1985, 1990, 1993b, 1997b,c, 1999, 2003; JENSEN & WELLS 1990; ICHIKAWA 1993; GOSLINER 1995; SWENNEN 1997, 2001; ORTEA & TEMPLADO 1988; ORTEA & ESPINOSA 1996, 2000, 2001, 2002; CABALLER et al. 2006; MUNIAN & ORTEA 1997). The present study has reviewed the existing information about sacoglossan distributions and analyzed the data for endemism and similarity among provinces. Although the data are biased with regards to collecting efforts of a few highly active scientists, several patterns have emerged from these analyses. The collection bias is most obvious in the areas where few other observations have been made, e.g. the Brazilian region where basically all collections have been made by the Marcuses. However, the species richness and endemism does not differ from that of other tropical regions with no collection bias. The same seems to be true for the Japanese fauna, where Baba has described most of the species. The sacoglossan fauna of the Indian sub-province has been studied by several taxonomists, but in this case endemism seems remarkably high. One reason for this is that descriptions have been published in local journals and therefore overlooked by other taxonomists.

Overall the distributions of sacoglossans correspond to the biogeographic regions and provinces identified by BRIGGS (1995). The exceptions have been pointed out above. Some of the provinces identified contain less than 10 species, which means that just one endemic species will

Table 8. Number of species and endemism of the tropical provinces and sub-provinces of the Indo-West Pacific region.

Province/ subprovince	# species	# endemics (%)
Western Indian Ocean	27	5 (19)
Red Sea	10	2 (20)
India <i>s.l.</i>	20	10 (50)
Indo-Malayan	48	9 (19)
Central Pacific	51	4 (8)
Ryukyu	33	10 (30)
Hawaii	25	6 (24)
S Pacific	33	6 (18)
Northeast Australia	28	2 (7)
+ Great Barrier Reef		
North and Northwestern Australia	23	4 (17)

yield more than 10% endemism. Obviously this cannot be used to make conclusions about their distinctiveness.

The records listed in existing literature do not usually have longitude and latitude, and many locality names are ambiguous or oblivious. Consequently it has not been possible to construct a "degree-by-degree" plot of species distributions for analysis. Nevertheless, the latitudinal decrease in species diversity from warm to cold temperate provinces is spectacular. Sacoglossans are dietary specialists and the majority of species feed on siphonaceous green algae, which are much more abundant and diverse in tropical and warm temperate waters. The cold temperate sacoglossan faunas of the North Atlantic as well as the Northeast Pacific Ocean are mostly eurythermic species with wide latitudinal distributions, extending well into neighboring warm water regions; indices of inclusion are over 60% (Tables 3b, 4b). This is also seen in Japan (Table 5b) and southern Africa between the tropical western Indian Ocean and the warm temperate southwestern Africa (Table 7b), but not nearly as pronounced in Australia (Table 6b). This apparently supports Rapoport's rule that species ranges in high latitudes are larger than in low latitudes. However, there are great differences between the provinces. The Northeast Pacific coldwater fauna is 43% of the tropical fauna, the Northeast Atlantic boreal fauna is 21% of the combined Lusitanian and Mediterranean fauna, and the Northwest Atlantic boreal fauna is only 12% of the Caribbean fauna.

The number of amphi-Atlantic species is relatively high, especially for the warm-water/tropical faunas (ORTEA et al. 1997; present study): 16 species (about 30%) occur in the Caribbean and the Lusitanian and/or Mediterranean provinces. However, amphi-Atlantic distributions for most of these species have only been recognized in recent years (TEMPLADO et al. 1990; ORTEA et al. 1988, 1998), and it cannot be excluded that human introductions are involved. Contrary to this, only one cold-temperate species, *Hermatea vancoverensis*, has an amphi-Pacific distribution. *Alderia modesta* has a circum-boreal/arctic distribution. *Elysiella pusilla* has recently been recorded from the Mexican-Panamanian province (BEHRENS & HERMOSILLO 2005). However, it cannot be excluded that this is a human introduction. The same is true for *Ercolania hoodleae*, a Japanese species, which has been recorded from California (BEHRENS 1991); this may also be a misidentification (TROWBRIDGE 2002). The so-called circum-tropical species, *Elysia ornata* and *Ercolania coerulea*, are conspicuously absent from tropical eastern Pacific; they extend from the Caribbean to the South and/or Central Pacific islands. *Lobiger souverbii*, which has been synonymized with several Indo-West Pacific species (KAY 1964; BABA 1974; GOSLINER 1987a; GOSLINER et al. 1996), and which occurs in the Caribbean as well

Table 9. Distribution of species of the Oxynoacea, Plakobranchoidea and Limapontioidea in provinces supported by present study.

Region	Oxynoacea	Plakobranchoidea	Limapontioidea
NE Atlantic	0	1	10
Lusitanian	4	10	26
Mediterranean	3	11	24
NW Atlantic	0	2	4
Caribbean	6	25	23
NE Pacif. warm + cold	0	1	9
Mexican-Panamanian	5	6	12
SE Pacific	4	2	3
Hawaii	7	13	5
Japan cold temp.	3	11	12
Japan warm temp.	9	18	14
Ryukyu Islands	9	15	11
Australia, tropical	9	20	14
Australia, warm + cold	8	10	11
W Indian Ocean	9	18	8
India, Sri Lanka, Maldi.	8	8	4
Indo-Malaya	5	29	14
Central Pacific	10	25	16
SW Pacific	12	17	4

as Mexican-Panamanian province (BEHRENS 1991; TROWBRIDGE 2002), may be the only truly circum-tropical species. Very few species are shared between the Caribbean and Mexican-Panamanian provinces, but there is probably a sister-taxon relationship between the East Pacific *Elysia diomedea* (formerly *Tridachiella diomedea*) and the Caribbean *E. crispata* (formerly *Tridachia crispata*) and *E. clarki*. Thus the Central American land bridge is as effective a geographical separation as the natural land barrier between the Red Sea and the Mediterranean. Also the few species occurring on both sides of the Isthmus of Panama may have arrived on the Pacific side as hitch-hikers on ships travelling through the Panama Canal.

The subdivision of the Indo-Polynesian province used in the present study may seem rather arbitrary. In many studies Papua New Guinea is considered a part of the Indo-Malayan sub-province, whereas it has been included in the South Pacific sub-province here. Only 14 species have been identified from Papua New Guinea, which is obviously an underrepresentation; GOSLINER (1992) gives 61 species for this area, but most are undescribed or have not been identified to species. Only sacoglossans that have been identified to species level can be used for calculating similarities between different sub-provinces. The 14 species are almost all widespread species found in the Indo-Malayan as well as the Central Pacific sub-provinces. The one species that does not occur in these sub-provinces, *Elysia expansa*, has its main distribution in tropical Australia. Geologically Papua New Guinea has been associated with the Australian continent (BRIGGS 1995).

The number of species decreases with longitude to the west of the Central Pacific islands, from 51 through 48 in the Indo-Malayan sub-province to 20 in the Indian sub-province and 27 in the Western Indian Ocean province. The majority of species in the latter province have widespread distributions, whereas less than half the species in the Indian sub-province belong in this category. A less pronounced longitudinal decrease is seen from 33 species in the South Pacific islands through 28 in northeastern Australia to 23 in northwestern Australia.

The number of species of the Central and South Pacific islands, including Ryukyu, is at least 50% higher than the number of species in tropical East Pacific. This is in spite of the fact that the combined area of these islands is relatively small. On the other hand the endemism of these islands is relatively low except in the Ryukyu Islands (Table 8). This means that insular isolation has not resulted in extensive speciation (or that subsequent dispersal has obscured such speciation). The high endemism of Ryukyu is almost entirely attributable to one fairly recent publication: ICHIKAWA (1993).

Looking at distribution within subordinal taxa of the Sacoglossa, it is obvious that the shelled suborder Oxynoacea, which all feed exclusively on algae of the genus *Caulerpa*, is restricted by the distribution of this alga. In the North Atlantic shelled sacoglossans occur in Bermuda, but there is a questionable record of *Ascobulla fragilis* from the northern part of Spain (see above). In the East Pacific shelled species are found in the Panamanian-

ian region, but also in the southernmost part of the Gulf of California. In the West Pacific shelled species occur in the warm temperate region. The only places where shelled species occur in cold temperate waters are Victoria, southeastern Australia and the west coast of central Japan. These places also have species diversity of *Canlerpa* rivaling many tropical and warm temperate provinces (DOUMENGE 1995; PRUD'HOMME VAN REINE et al. 1996). The two non-shelled superfamilies, Plakobranchoidea and Limapontioidea, occur in all provinces where sacoglossans are found.

At the family level, the bivalved Juliidae are absent from the eastern Atlantic. Since they are common as fossils from this region, this must be due to extinction. Juliidae and Volvatellidae are rare in the Atlantic and eastern Pacific. On the other hand, Limapontioidea are highly diverse in the northeastern Atlantic and Mediterranean (Table 9). The least studied areas have the fewest Limapontioidea. This is probably due to the very small size (<10 mm) of most of these species. In tropical regions the number of Plakobranchoidea is always higher than the number of species in the other superfamilies. Plakobranchoidea are most diverse in the Caribbean region and in the Indo-Malayan and Central Pacific islands sub-provinces; tropical Australia also has a high proportion of this superfamily (Table 9).

Only rarely are more than two or three species of one genus found in one area. However, for the genera *Elysia* and *Thuridilla* many places have more than 5 species, and in a few cases more than 10 species may co-occur, though on different food algae. Whether this is due to sym- or parapatric speciation or dispersal following allopatric speciation, as suggested by GOSLINER (1995), cannot be deduced from existing information. Phylogenetic analysis, preferably including molecular data should be applied to these genera¹⁾.

Some genera are restricted to one or a few neighboring biogeographic provinces. *Limapontia* only occurs in the Northeast Atlantic and Mediterranean region. This may also be the case for the genus *Calliopaea*, though the Japanese *Stiliger pusillus* has been tentatively assigned to this genus (BABA & HAMATANI 1970). The monotypic *Platyhedyle* only occurs in the Mediterranean (WAWRA 1979), whereas the genus *Gascoignella*, which has been assigned to the same family, Platyhedylidae (JENSEN 1996a), seems restricted to the South China Sea (JENSEN 1985; SWENNEN 2001). The monotypic genus *Olea* is restricted to the Northeast Pacific (TROWBRIDGE 2002). The genus *Bosellia* may have its natural distribution restricted to tropical and warm temperate Atlantic and Mediterranean (MARCUS 1973; FERNANDEZ-OVIES & ORTEA 1986), and the monotypic genus *Sohgenia* has only been found in the Central Pacific islands (HAMATANI 1991). Common to all

of the above genera is that they have morphological characters that appear to be reduced compared to other genera in the same superfamilies; they either lack or have very reduced rhinophores, cerata or parapodia (or a combination of these). The genera *Roburriella*, *Plakobranchus* and *Pattyclaya* are Indo-west Pacific endemics, the former furthermore restricted to southern Australia, and *Julia* and *Elysia* also occur in the eastern Pacific. Within the more speciose genera, many have few (1-3) representatives in the Atlantic Ocean, zero in the East Pacific, and the remaining species are Indo-West Pacific. This is seen in *Volvatella*, *Thuridilla* and *Costasiella*, indicating that the Atlantic fauna is a Tethyan relict fauna with little subsequent speciation, except in the genus *Elysia* in the Caribbean. However, the high number of limapontioid species seems to contradict this. It is possible that the lack of plakobranchoids first spurred a burst of speciation of limapontioids in the East Atlantic/Mediterranean, whereas the speciation in Caribbean *Elysia* may be a more recent phenomenon.

The present study is a preliminary analysis of biogeographic affinities of sacoglossan opisthobranchs. Nevertheless several patterns have been identified, which should be further investigated using phylogenetic analyses. Also, some obvious gaps in the existing knowledge as well as conflicting and/or questionable records have been identified. There is an urgent need to describe the undescribed species known from the Indo-West Pacific region and to collect sacoglossans from poorly studied regions. Finally, several new hypotheses emerge from the present analyses, which should be tested in the future: the genus *Elysia* has speciated within the Caribbean; warm temperate Australia and Japan are centers of speciation; the appearance of several Caribbean species in the Canary Islands is due to human activities.

Acknowledgements. I would like to thank Heike Wägele (Bonn) and the organizing committee for organizing the Second International Workshop on Opisthobranchia.

REFERENCES

- BABA, K. 1935. The fauna of Akkeshi Bay. 1. Opisthobranchia. Journal of the Faculty of Science, Hokkaido Imperial University, series 6, Zoology 4: 115–125, plates 7–8.
- BABA, K. 1936. Opisthobranchia of the Ryukyu (Okinawa) Islands. Journal of the Department of Agriculture, Kyushu Imperial University 5: 1–49, plates 1–3.
- BABA, K. 1949. Opisthobranchia of Sagami Bay. Collected by His Majesty the Emperor of Japan. Tokyo: Iwanami Shoten.
- BABA, K. 1952a. Record of an ascoglossan mollusk, *Oxynoe viridis* (Pease) from Sagami Bay, Japan. Venus (Japanese Journal of Malacology) 17(2): 77–80.
- BABA, K. 1952b. Record of a rare sacoglossan mollusk, *Lobiger* (*Lobiger*) *sagamiensis* n.sp. from Sagami Bay, Japan. Zoological Magazine (Dobutsugaku Zasshi) 61(11): 337–338.

¹⁾After the acceptance of this paper, the following article was published: BASS, A.L. & KARL, S.A. 2006. Molecular phylogenetic analysis of genera in the family Plakobranchoidea (Mollusca: Opisthobranchia: Sacoglossa). In: Contemporary studies into the systematics and evolution of opisthobranch molluscs (eds. G. BRODIE, S. FAHEY & F.E. WELLS). Records of the Western Australian Museum, Supplement 69, Western Australian Museum, Perth, WA.

- BABA, K. 1955. Opisthobranchia of Sagami Bay, Supplement. Collected by His Majesty The Emperor of Japan. Tokyo: Iwanami Shoten, 59pp, 20 colour plates.
- BABA, K. 1957. The species of the genus *Elysia* from Japan. Publications of the Seto Marine Biological Laboratory 6: 69–74, plates 3–4.
- BABA, K. 1959. The family Stiligeridae from Japan (Opisthobranchia - Sacoglossa). Publications of the Seto Marine Biological Laboratory 7: 327–334, plates 27–28.
- BABA, K. 1966. Gross anatomy of the specimens of the shelled sacoglossan *Volvatella* (= *Arthessa*) collected from Okino-Erubu Island, southern Kyushu, Japan (Nudibranchia). Publications of the Seto Marine Biological Laboratory 14: 197–205, plates 7–10.
- BABA, K. 1968. A revised description of *Alderia nigra* Baba, 1937, type species of *Alderiopsis* n.g. from Japan (Opisthobranchia - Sacoglossa). Bijdragen tot de Dierkunde 38: 5–11.
- BABA, K. 1974. Some comments on *Lobiger souverbii* Fischer, 1856, re-identified, of Japan. The Veliger 16: 253–257.
- BABA, K. & HAMATANI, I. 1970. The anatomy of *Ercolania boodleeae* (Baba, 1938) from Seto, Kii, Middle Japan (Opisthobranchia: Sacoglossa). Publications of the Seto Marine Biological Laboratory 18: 215–222, plates 5–7.
- BARASH, A. & DANIN, Z. 1971. Opisthobranchia (Mollusca) from the Mediterranean waters of Israel. Israel Journal of Zoology 20: 151–200.
- BEHRENS, D.W. 1991. Pacific coast nudibranchs. Second edition. Sea Challengers, Monterey, California.
- BEHRENS, D.W. & HERMOSILLO, A. 2005. Eastern Pacific Nudibranchs. A Guide to the Opisthobranchs from Alaska to Central America. Sea Challengers, Monterey, California, 137pp.
- BERGH, R. 1871. Malacologische Untersuchungen. Reisen im Archipel der Philippinen von C. Semper 2(2): 49–118, plates 9–16.
- BERGH, R. 1872. Malacologische Untersuchungen.. Reisen im Archipel der Philippinen von C. Semper 2(3–4): 137–203, plates 17–24.
- BERGH, R. 1888. Malacologische Untersuchungen. Reisen im Archipel der Philippinen von C. Semper 3(16): 755–814, plates 77–81.
- BERGH, R. 1905. Die Opisthobranchiata der Siboga Expedition. Brill, Leiden.
- BLEAKNEY, J.S. 1989. Morphological variation in the radula of *Placida dendritica* (Alder & Hancock, 1843) (Opisthobranchia: Ascoglossa/Sacoglossa) from Atlantic and Pacific populations. The Veliger 32: 171–181.
- BLEAKNEY, J.S. 1996. Sea Slugs of Atlantic Canada and the Gulf of Maine. The Nova Scotia Museum Field Guide Series. Nimbus Publishing and The Nova Scotia Museum: Halifax, 215pp.
- BOETTGER, C. 1963. Gastropoden mit zwei Schalenklappen. Zoologischer Anzeiger, Supplement 26: 403–439.
- BOONE, C.E. 1982. More on *Elysia* in Texas. Texas Conchologist 18: 29–37.
- BOUCHET, P. 1984. Les Elysiidae de Méditerranée (Gastropoda, Opisthobranchiata). Annales de l'Institut Océanographique, Paris 60: 19–28.
- BRATTEGARD, T. & HOLTHE, T. 2001. Distribution of marine, benthic macroorganisms in Norway. A tabulated catalogue. Update of Utredning for DN 1997–1. Directorate for Nature Management, Trondheim, Norway. Research Report 2001–3, 409 pp.
- BRIGGS, J.C. 1995. Global Biogeography. Developments in Palaeontology and Stratigraphy 14. Elsevier, 454pp.
- BRIGGS, J.C. 2005. The marine East Indies: diversity and speciation. Journal of Biogeography 32: 1517–1522.
- BRODIE, G.D. & BRODIE, J.E. 1990. A checklist of the opisthobranch molluscs of Fiji. Journal of the Malacological Society of Australia 11: 53–63.
- BURN, R. 1958. Further Victorian Opisthobranchia. Journal of the Malacological Society of Australia 1(2): 20–36.
- BURN, R. 1960. Occurrence of bivalve gastropods along the coast-line of New South Wales. Nature 188: 680–681.
- BURN, R. 1965. Rediscovery and taxonomy of *Edentellina typica* Gatliff & Gabriel. Nature 206: 735–736.
- BURN, R. 1966a. The opisthobranchs of a Caulerpa microfauna from Fiji. Proceedings of the Malacological Society of London 37: 45–65.
- BURN, R. 1966b. Some opisthobranchs from southern Queensland. Journal of the Malacological Society of Australia 1(9): 96–109.
- BURN, R. 1973. *Limapontia* in Australia. Australian Shell News 4: 2.
- BURN, R. 1974. Notes on some benthonic opisthobranchs from Port Phillip Bay, Victoria. Journal of the Malacological Society of Australia 3: 43–57.
- BURN, R. 1998. Order Sacoglossa. Pp. 961–974 in: BEESLEY, P.L., ROSS, G.J.B. & WELLS, A. (eds.) Mollusca: The Southern Synthesis. Fauna of Australia. Vol. 5. CSIRO Publishing: Melbourne, Part B, viii 565–1234.
- BURN, R. 2006. A checklist and bibliography of the Opisthobranchia (Mollusca: Gastropoda) of Victoria and the Bass Strait area, south-eastern Australia. Museum Victoria Science Reports 10: 1–42.
- CABALLER, M., ORTEA, J. & ESPINOSA, J. 2006. Descripción de una nueva especie de *Alderiopsis* Baba, 1968. In: Moluscos marinos de la península de Guanahacabibes, Pinar del Río, Cuba, con la descripción de nuevos taxones. Avicennia 18: 57–59.
- CARLSON C.H. & HOFF, P.-J. 1978. The identifiable *Elysia* from Guam (Elysiidae, Sacoglossa, Opisthobranchia). Micronesica 14: 89–113.
- CARLSON, C. & HOFF, P.-J. 2003. The opisthobranchs of the Mariana Islands. Micronesica 35–36: 272–295.
- CATTANEO-VIETTI, R. & THOMPSON, T.E. 1989. Mediterranean opisthobranch molluscs: a zoogeographic approach. Bolletino Malacologico 25: 183–204.
- CERVERA, J.L., CALADO, G., GAVAIÁ, C., MALAQUIAS, M.A.E., TEMPLADO, J., BALLESTEROS, M., GARCIA-GÓMEZ, J.C. & MEGINA, C. 2006. An annotated and updated checklist of the opisthobranchs (Mollusca: Gastropoda) from Spain and Portugal (including islands and archipelagos). Boletín Instituto Español de Oceanografía 20(1–4): 1–122.
- CERVERA, J.L., GARCIA-GÓMEZ, J.C. & ORTEA, J.A. 1991. Una nueva especie del género *Hermaea* (Gastropoda: Opisthobranchia: Sacoglossa) y redescription de dos raros sacoglossos de la malacofauna Europea. Iberus 8: 215–224.
- CERVERA, J.L. & LOPEZ-GONZÁLEZ, P.J. 1996. New records of two uncommon sacoglossans (Gastropoda: Opisthobranchia) from the coasts of the Iberian Peninsula. The Veliger 39: 93–95.
- CERVERA, J.L., TEMPLADO, J., GARCIA-GÓMEZ, J.C., BALLESTEROS, M., ORTEA, J.A., GARCIA, F.J., ROS, J. & LUQUE, A.A. 1988. Catalogo actualizado y comentado de los opisthobranchios (Mollusca, Gastropoda) de la península Iberica, Baleares y Canarias, con algunas referencias a Ceuta y la Isla de Alboran. Iberus, Supplement 1: 7–84.
- CHAPMAN, J.W. & CARLTON, J.T. 1991. A test of criteria for introduced species: the global invasion by the isopod *Synidotea laevidorsalis* (Miers, 1881). Journal of Crustacean Biology 11: 386–400.

- CHERNYSHEV, A. 2005. *Stiliger akkeshiensis* from Kuril Islands. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=14612>) (last access 28th of August 2007).
- CHERNYSHEV, A.V. & CHABAN, E.M. 2005. *Alderia modesta* from eastern Russia. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=13381>) (last access 28th of August 2007).
- CLARK, K.B. 1975. Nudibranch life cycles in the northwest Atlantic and their relationship to the ecology of fouling communities. *Helgoländer Wissenschaftliche Meeresuntersuchungen* **27**: 28–69.
- CLARK, K.B. 1982. A new *Vohatella* (Mollusca: Ascoglossa) from Bermuda, with comments on the genus. *Bulletin of Marine Science* **32**: 112–20.
- CLARK, K.B. 1984. New records and synonymies of Bermuda opisthobranchs (Gastropoda). *Nautilus* **98**: 85–97.
- CLARK, K.B. & DE-FRESE, D. 1987. Population ecology of Caribbean Ascoglossa (Mollusca: Opisthobranchia): A study of specialized algal herbivores. *American Malacological Bulletin* **5**: 259–280.
- DEBELIUS, H. 1996. Nudibranchs and Sea Snails. Indo-Pacific Field Guide. IKAN-Underwasserarchiv, Frankfurt, 321pp.
- DOUMENGE, F. 1995. Quelques réflexions sur les algues caulerpes. *Biologia Marina Mediterranea* **2**: 613–633.
- EDMUNDS, M. 1963. *Berthelinia caribbea* n.sp., a bivalved gastropod from the West Atlantic. *Journal of the Linnean Society of London, Zoology* **44**: 731–739.
- EKMAN, S. 1953. Zoogeography of the Sea. Sidgwick & Jackson.
- ELIOT, C. 1899. Notes on tectibranchs and naked mollusks from Samoa. *Proceedings of the Academy of Natural Sciences of Philadelphia* **1899**: 512–523, pl. 19.
- ELIOT, C. 1903. On some nudibranchs from East Africa and Zanzibar. Part II. *Proceedings of the Zoological Society of London* **1903**(1): 250–257.
- ELIOT, C. 1904. On some nudibranchs from East Africa and Zanzibar. Part VI. *Proceedings of the Zoological Society of London* **1904**(2): 268–298.
- ELIOT, C. 1906. On the nudibranchs of Southern India and Ceylon with special reference to the drawings by Kelaart and collections belonging to Alder and Hancock preserved in the Hancock Museum at Newcastle-on-Tyne. *Proceedings of the Zoological Society of London* **1906**: 636–691.
- ELIOT, C. 1908. Reports on the marine biology of the Sudanese Red Sea. XI. Notes on a collection of nudibranchs from the Red Sea. *Journal of the Linnean Society* **31**: 86–122.
- ELIOT, C. 1917. Mollusca Nudibranchiata (Ascoglossa). Pp. 179–182 in: ANNANDALE, N. (ed.) *Zoological Results of a Tour in the Far East, Part III. Memoirs of the Asiatic Society of Bengal* **6**.
- ENGEL, H., GEERTS, S.J. & VAN REGTEREN-ALTENA, C.O. 1940. *Alderia modesta* (Lovén) and *Limapontia depressa* Alder & Hancock in the brackish waters of the Dutch coast. *Basteria* **5**: 6–34.
- EVERTSEN, J. & BAKKEN, T. 2002. Heterobranchia (Mollusca, Opisthobranchia) from northern Norway, with notes on ecology and distribution. *Fauna Norvegica* **22**: 15–22.
- FERNANDEZ-OVIES, C.L. & ORTEA, J. 1986. Descripción de una nueva especie de *Bosellia* Trinchese, 1890 (Mollusca: Opisthobranchia: Ascoglossa) de las Islas Canarias. *Iberus* **6**: 101–106.
- GODDARD, J.H.R. & FOSTER, N.R. 2002. Range extensions of sacoglossan and nudibranch mollusks (Gastropoda: Opisthobranchia) to Alaska. *The Veliger* **45**(4): 331–336.
- GOLIKOV, A.N. 1989. Arctic Ocean gastropods, prosobranchs. Pp. 325–340. In: HERMAN, Y. (ed.) *The Arctic Seas: Climatology, Oceanography, Geology, and Biology*. Van Nostrand Reinhold, New York. 888pp.
- GOSLINER, T.M. 1987a. Biogeography of the opisthobranch gastropod fauna of southern Africa. *American Malacological Bulletin* **5**: 243–258.
- GOSLINER, T.M. 1987b. Nudibranchs of southern Africa. A guide to opisthobranch molluscs of southern Africa. Sea Challengers and Jeff Hamann, Monterey and El Cajon, California. 136 pp.
- GOSLINER, T.M. 1992. Biodiversity of tropical opisthobranch gastropod faunas. *Proceedings of the Seventh International Coral Reef Symposium* **2**: 702–709.
- GOSLINER, T.M. 1995. The genus *Thuridilla* (Opisthobranchia: Elysiidae) from the tropical Indo-Pacific, with a revision of the phylogeny and systematics of the Elysiidae. *Proceedings of the California Academy of Sciences* **49**: 1–54.
- GOSLINER, T.M. 1998. Name changes. Update for Gosliner 1987: Nudibranchs of Southern Africa. *Australasian nudibranch news* No. **3** (Available on: <http://slugsite.tierranet.com/news/annws3.pdf>)
- GOSLINER, T.M., BEHRENS, D.W. & WILLIAMS, G.C. 1996. Coral reef animals of the Indo-Pacific. Sea Challengers, Monterey, California, 314pp.
- GOSLINER, T.M. & DRAHEIM, R. 1996. Indo-Pacific opisthobranch gastropod biogeography: how do we know what we don't know? *American Malacological Bulletin* **12**: 37–43.
- GOULD, A.A. & BINNEY, W.G. 1870. Report on the invertebrates of Massachusetts. 524pp. + 27 plates.
- GULLIKSEN, B., PALERUD, R., BRATTEGARD, T. & SNELI, J.-A. 1999. Distribution of marine benthic macro-organisms at Svalbard (including Bear Island) and Jan Mayen. Research Report for DN 1999-4. Directorate for Nature Management, Trondheim, Norway. 148pp.
- HAMATANI, I. 1968. A new species of *Elysia* from Kii, Japan (Opisthobranchia - Sacoglossa). *Publications of the Seto Marine Biological Laboratory* **16**: 51–54.
- HAMATANI, I. 1969. A new species of the rare shelled sacoglossan genus *Cylindrobulla* from Middle Japan. *Publications of the Seto Marine Biological Laboratory* **17**: 171–174, pls. 5–6.
- HAMATANI, I. 1971. A new species of *Cylindrobulla*, sacoglossan opisthobranch, from California; with a comparison with *C. japonica* Hamatani, 1969. *Publications of the Seto Marine Biological Laboratory* **19**: 111–117, plates 6–7.
- HAMATANI, I. 1972. A new species of *Vohatella* Pease 1860, found in the "Caulerpan microfauna" in the province of Kii, middle Japan (Opisthobranchia, Sacoglossa). *Publications of the Seto Marine Biological Laboratory* **21**: 13–20, pls. 2–3.
- HAMATANI, I. 1976a. Preliminary account of a new species of *Vohatella* Pease, 1860, *V. viridis* sp. nov., found in the Caulerpan microfauna in Japan (Opisthobranchia: Ascoglossa). *Publications of the Seto Marine Biological Laboratory* **22**: 371–376, plate 6.
- HAMATANI, I. 1976b. A new species of *Cyerce* Bergh, 1871, *C. kikutarobabai* from Yoron Islands (Opisthobranchia: Sacoglossa). *Publications of the Seto Marine Biological Laboratory* **23**: 283–288.
- HAMATANI, I. 1980. On the species of the genus *Oxynoe* Rafinesque, 1819 from Japan, inclusive of a new species (Opisthobranchia: Ascoglossa). *Publications of the Seto Marine Biological Laboratory* **25**: 349–360, plates 1–4.
- HAMATANI, I. 1991. *Sohgenia palauensis* n.gen. & sp., a new ascoglossan opisthobranch from the Palau Islands collected by

- the R/V Sohgen-Mar. Venus (Japanese Journal of Malacology) **50**: 85–92.
- HAMATANI, I. 1994. A new species of *Mourgona* Marcus & Marcus, 1970, *M. osumi* n.sp. (Opisthobranchia: Ascoglossa), found in acetabularian microfauna from Amami-Oshima Island, Southwestern Japan. Venus (Japanese Journal of Malacology) **53**: 21–27.
- HELLER, J. & THOMPSON, T.E. 1983. Opisthobranch molluscs of the Sudanese Red Sea. Zoological Journal of the Linnean Society **78**: 317–348.
- HIRANO, Y.J., HIRANO, Y.M. & TROWBRIDGE, C.D. 2006. Record of a common but cryptic sacoglossan, *Placida daguilarensis* Jensen 1990, from Japan. Memoirs of the National Science Museum, Tokyo **40**: 279–290.
- CHIKAWA, M. 1999. Sacoglossa (Opisthobranchia) from the Ryukyu Islands. Publications of the Seto Marine Biological Laboratory **36**: 119–139.
- IMAMOTO, J. 2004. *Bosellia mimetica?* from Japan. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=11858>) (last access 28th of August 2007).
- JENSEN, K.R. 1982. Occurrence of *Elysia serca* Marcus in Florida, with notes on the synonymy of the species. Journal of Conchology **31**: 87–94.
- JENSEN, K.R. 1985. Annotated checklist of Hong Kong Ascoglossa (Mollusca: Opisthobranchia), with descriptions of four new species. Pp. 77–107 in: MORTON, B. & DUDGEON, D. (eds.) Proceedings of the Second International Workshop on the Malacofauna Hong Kong and Southern China, Hong Kong 1983, 1. Hong Kong University Press, Hong Kong.
- JENSEN, K.R. 1990. Three new species of Ascoglossa (Mollusca, Opisthobranchia) from Hong Kong, and a description of the internal anatomy of *Costasiella pallida* Jensen, 1985. Pp. 419–432 in: MORTON, B. (ed.) The marine flora and fauna of Hong Kong and southern China II. Proceedings of the Second International Marine Biological Workshop: The Marine Flora and Fauna of Hong Kong and Southern China, Hong Kong, 2–24 April 1986. Hong Kong University Press, Hong Kong.
- JENSEN, K.R. 1993a. Morphological adaptations and plasticity of radular teeth of the Sacoglossa (=Ascoglossa) (Mollusca: Opisthobranchia) in relation to their food plants. Biological Journal of the Linnean Society **48**: 135–155.
- JENSEN, K.R. 1993b. Sacoglossa (Mollusca, Opisthobranchia) from Rottnest Island and central Western Australia. Pp. 207–253 in: WELLS, F.E., WALKER, D.I., KIRKMAN, H. & LETHBRIDGE, R. (eds.) Proceedings of the Fifth International Marine Biological Workshop: The Marine Flora and Fauna of Rottnest Island, Western Australia. Western Australian Museum, Perth.
- JENSEN, K.R. 1995. Anatomy and biology of *Aphysiopsis formosa* Pruvot-Fol (Mollusca, Opisthobranchia, Sacoglossa) from the Azores. Açoreana, Supplement 1995: 217–230.
- JENSEN, K.R. 1996a. Phylogenetic systematics and classification of the Sacoglossa (Mollusca, Gastropoda, Opisthobranchia). Philosophical Transactions of the Royal Society B **351**: 91–122.
- JENSEN, K.R. 1996b. The Diaphanidae as a possible sister group of the Sacoglossa (Gastropoda, Opisthobranchia). Pp. 231–247 in: TAYLOR, J. (ed.) Origin and evolutionary radiation of the Mollusca. Oxford University Press, Oxford.
- JENSEN, K.R. 1997a. Evolution of the Sacoglossa (Mollusca, Opisthobranchia) and the ecological associations with their food plants. Evolutionary Ecology **11**: 301–335.
- JENSEN, K.R. 1997b. Sacoglossa (Mollusca, Opisthobranchia) from the Darwin Harbour area, Northern Territory, Australia. Pp. 163–186 in: HANLEY, J.R., CASWELL, G., MEGIRIAN, D. & LARSON, H.K. (eds.) Proceedings of the Sixth International Marine Biological Workshop: The Marine Flora and Fauna of Darwin, Northern Territory, Australia. Darwin, Australia: Museums and Art Galleries of the Northern Territory and the Australian Marine Sciences Association.
- JENSEN, K.R. 1997c. Sacoglossa (Mollusca, Opisthobranchia) from the Houtman Abrolhos Island and central Western Australia. Pp. 307–333 in: WELLS, F.E. (ed.) Proceedings of the Seventh International Marine Biological Workshop: The Marine Flora and Fauna of the Houtman Abrolhos Islands, Western Australia. Western Australian Museum, Perth.
- JENSEN, K.R. 1998a. Zoogeographic affinities of Hong Kong Opisthobranchia (Mollusca, Gastropoda). Pp. 43–55 in: MORTON, B. (ed.) The Marine Biology of the South China Sea. Proceedings of the Third International Conference on the Marine Biology of the South China Sea, Hong Kong, 28 October–1 November 1996. Hong Kong: Hong Kong Univer. Press.
- JENSEN, K.R. 1998b. Anatomy of some opisthobranch molluscs from Phuket, Thailand, with a list of Opisthobranchia recorded from Thailand. Phuket Marine Biological Center Special Publication **18**(2): 243–262.
- JENSEN, K.R. 1999. A new species of Sacoglossa (Mollusca, Opisthobranchia) from Rottnest Island, Western Australia. Pp. 377–383 in: WALKER, D.I. & WELLS, F.E. (eds.) The Seagrass Flora and Fauna of Rottnest Island, Western Australia. Western Australian Museum: Perth.
- JENSEN, K.R. 2003. Distributions, diets and reproduction of Hong Kong Sacoglossa (Mollusca: Opisthobranchia): a summary of data, 1980–2001. Pp. 347–365 in: MORTON, B. (ed.) Perspectives on Marine Environmental Change in Hong Kong and Southern China, 1977–2001. Proceedings of an International Workshops Reunion Conference, Hong Kong 2001, Hong Kong University Press: Hong Kong.
- JENSEN, K.R. & CLARK, K.B. 1983. Annotated checklist of Florida ascoglossan Opisthobranchia. Nautilus **97**: 1–13.
- JENSEN, K.R. & WELLS, F.E. 1990. Sacoglossa (=Ascoglossa) (Mollusca, Opisthobranchia) from southern Western Australia. Pp. 297–331 in: WELLS, F.E., WALKER, D.I., KIRKMAN, H. & LETHBRIDGE, R. (eds.) Proceedings of the Third International Marine Biological Workshop: The Marine Flora and Fauna of Albany, Western Australia I. Western Australian Museum, Perth.
- JOHNSON, S. & BOUCHER, L.M. 1983. Notes on some Opisthobranchia (Mollusca: Gastropoda) from the Marshall Islands, including 57 new records. Pacific Science **37**: 251–291.
- KAWAGUTI, S. & BABA, K. 1959. A preliminary note on a two-valved sacoglossan gastropod, *Tamanovalva limax* n.gen., n.sp., from Tamano, Japan. Biological Journal Okayama University **5**: 177–184.
- KAY, E.A. 1964. A new species of *Berthelinia* and its associated sacoglossans in the Hawaiian Islands. Proceedings of the Malacological Society of London **36**: 191–197.
- KAY, E.A. 1967. The composition and relationships of marine molluscan fauna of the Hawaiian islands. Venus (Japanese Journal of Malacology) **25**(3–4): 94–104.
- KAY, E.A. 1968. A review of the bivalved gastropods and a discussion of evolution within the Sacoglossa. Symposia of the Zoological Society of London **22**: 109–134.
- KEEN, A. M. & SMITH, A.G. 1961. West American species of the gastropod genus *Berthelinia*. Proceedings of the California Academy of Science **30**: 47–66.
- KELAART, E.F. 1858. Descriptions of new and little-known species of Ceylon nudibranchiate molluscs and zoophytes. Journal of the Royal Asiatic Society, Ceylon Branch **3**(1): 84–139.

- KOH, D.B. 2002a. *Elysia amakusana* from South Korea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=8250>) (last access 12th of August 2007).
- KOH, D.B. 2002b. *Elysia ornata* from South Korea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=8256>) (last access 12th of August 2007).
- KOH, D.B. 2003. *Hermatea bifida?* From Korea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=11522>) (last access 12th of August 2007).
- KOH, D.B. 2005a. *Aplysiopsis nigra* (Baba, 1949) from Sth Korea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=13403>) (last access 12th of August 2007).
- KOH, D.B. 2005b. *Elysia obtusa* from South Korea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=13964>) (last access 12th of August 2007).
- KOH, D.B. 2005c. *Elysia atroviridis* from Sth Korea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=14126>) (last access 12th of August 2007).
- KRUG, P.J., ELLINGSON, R.A., BURTON, R.A. & VALDÉS, A. 2007. A new poecilogenous species of sea slug (Opisthobranchia: Sacoglossa) from California: Comparison with the planktotrophic congener *Alderia modesta* (Lovén, 1844). *Journal of Molluscan Studies* **73**: 29–38.
- KURAKIN, A. 2002. Opisthobranch from the Black Sea. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=8300>).
- LE RENARD, J., SABELLI, B. & TAVIANI, M. 1996. On *Candina* (Sacoglossa: Juliidae), a new fossil genus of bivalved gastropods. *Journal of Paleontology* **70**(2): 230–235.
- LEAL, J.H. & BOUCHET, P. 1991. Distribution patterns and dispersal of prosobranch gastropods along a seamount chain in the Atlantic Ocean. *Journal of the Marine Biological Association of the United Kingdom* **71**: 11–25.
- LEMICHE, H. & THOMPSON, T.E. 1974. Three opisthobranch gastropods new to the British Fauna. *Proceedings of the Malacological Society of London* **41**: 185–193.
- LIN, G. 1986. Additions to the Opisthobranchia fauna of the Hainan and Xisha Islands of China. *Studia Marina Sinica* **26**: 117–128, plates 1–2.
- LIN, G. 1990. Opisthobranchia fauna of the Hainan Island, China. *Bulletin of Marine Science* **47**: 134–138.
- MACNAE, W. 1954. On four sacoglossans new to South Africa. *Annals of the Natal Museum* **13**: 51–64, plate 3.
- MALACUIAS, M.A.E. & CALADO, G.J.P. 1997. The malacological fauna of Salvage Islands. 1. Opisthobranch molluscs. *Boletim Muscu Municipal Funchal* **49**(281): 149–170.
- MARCUS, E. 1955. Opisthobranchia from Brazil. *Boletins da Faculdade de Filosofia da Universidade de Sao Paulo, Zoologia* **20**: 89–262, plates 1–30.
- MARCUS, E. 1957. On Opisthobranchia from Brazil (2). *Journal of the Linnean Society, Zoology* **43**: 390–486.
- MARCUS, E. 1959. Lamellariacea und Opisthobranchia. Reports of the Lund University Chile Expedition 1948–49. *Lunds Universitets Årsskrift* **55**: 1–133.
- MARCUS, E. 1965. Some opisthobranchs from Micronesia. *Malacologia* **3**: 263–286.
- MARCUS, E. & MARCUS, E. 1966. The R/V Pilsbry deep-sea biological expedition to the Gulf of Guinea, 1964–65: 9 Opisthobranchs from tropical West Africa. *Studies in Tropical Oceanography, Miami* **4**: 152–208.
- MARCUS, E. & MARCUS, E. 1967. Tropical American opisthobranchs. *Studies in Tropical Oceanography, Miami* **6**: 3–137.
- MARCUS, E. & MARCUS, E. 1970. Opisthobranchs from Curaçao and faunistically related regions. *Studies on the Fauna of Curaçao and other Caribbean Islands* **33**: 1–129.
- MARCUS, E. DEB.-R. 1972a. Notes on some opisthobranch gastropods from the Chesapeake Bay. *Chesapeake Science* **13**: 300–317.
- MARCUS, E. DEB.-R. 1972b. On some opisthobranchs from Florida. *Bulletin of Marine Science* **22**: 284–308.
- MARCUS, E. DEB.-R. 1973. On the genus *Bosellia* (Mollusca: Gastropoda: Ascoglossa). *Bulletin of Marine Science* **23**: 811–823.
- MARCUS, E. DEB.-R. 1977. An annotated checklist of Western Atlantic warm water opisthobranchs. *Journal of Molluscan Studies, Supplement* **4**: 1–21.
- MARCUS, E. DEB.-R. 1978. On a new species of *Bosellia*. *Boletim de Zoologia, Universidade de Sao Paulo* **3**: 1–6.
- MARCUS, E. DEB.-R. 1980. Review of western Atlantic Elysiidae (Opisthobranchia Ascoglossa) with a description of a new *Elysia* species. *Bulletin of Marine Science* **30**: 54–79.
- MARCUS, E. DEB.-R. & HUGHES, H.P.I. 1974. Opisthobranch molluscs from Barbados. *Bulletin of Marine Science* **24**: 498–532.
- MARSHALL, J.G. & WILLAN, R.C. 1999. Nudibranchs of Heron Island, Great Barrier Reef. A survey of the Opisthobranchia (Sea Slugs) of Heron and Wistari Reefs. *Backhuys Publishers, Leiden*, 257pp.
- MARTYNOV, A.V., KORSHUNOVA, T.A. & SAVINKIN, O.V. 2006. Shallow-water opisthobranch mollusc of the Murman coast of the Barents Sea, with new distributional data and remarks on biology. *Ruthenica* **16**(1–2): 59–72.
- MIKKELSEN, P.M. 1996. The evolutionary relationships of Cephalaspidea s.l. (Gastropoda: Opisthobranchia): A phylogenetic analysis. *Malacologia* **37**: 375–442.
- MIKKELSEN, P.M. 1998. *Cylindrobulla* and *Ascobulla* in the western Atlantic (Gastropoda, Opisthobranchia, Sacoglossa): Systematic review, description of a new species, and phylogenetic reanalysis. *Zoologica Scripta* **27**: 49–71.
- MILLER, M.C. 1969. The habits and habitats of the opisthobranch molluscs of the British Solomon Islands. *Philosophical Transactions of the Royal Society B* **255**: 541–548.
- MUNIAN, C. & ORTEA, J. 1997. First record of a sacoglossan (= ascoglossan, Opisthobranchia) from Patagonia (Argentina): Description of a new species of genus *Elysia* Risso, 1818. *The Veliger* **40**: 29–37.
- MURINA, V.V. & ARTEMJEVA, J.N. 1997. Phenology of molluscan pelagic larvae from the southern coastal waters of Crimea. *Scientia Marina* **61** (Supplement 2): 55–58.
- NEVILL, G. & NEVILL, H. 1869. On some new marine Gastropoda from the southern province of Ceylon. *Journal of Asiatic Society, Bengal* **38**: 65–69.
- O'DONOGHUE, C. 1929. Report on the Nudibranchiata. Zoological results of the Cambridge Expedition to the Suez Canal 1927. *Transactions of the Zoological Society of London* **22**: 713–841.
- O'DONOGHUE, C.H. 1932. Kelaart's work on the Nudibranchiata of Ceylon. *Proceedings of the Malacological Society of London* **20**: 221–226.
- ORTEA, J. 1981. Moluscos opisthobranchios de las Islas Canarias. 1. Parte: Ascoglossos. *Boletim do Instituto España Oceanográfico* **6**: 180–199.
- ORTEA, J., BACALLADO, J.J. & SANCHEZ, J.M.P. 1990. *Aplysiopsis formosa* Pruvot-Fol, 1953 (Mollusca, Opisthobranchia, Ascoglossa) in the Canary Islands. *Lavori della Società Italiana di Malacologia, Napoli* **23**: 281–285.

- ORTEA, J. & ESPINOSA, J. 1996. Descripción de una nueva especie del género *Elysia* Risso, 1818 (Opisthobranchia: Sacoglossa) recolectada en Puerto Morelos, Mexico. *Avicennia* **4-5**: 115–119.
- ORTEA, J. & ESPINOSA, J. 2000. Nueva especie del género *Thuridilla* Bergh, 1872 (Mollusca: Sacoglossa) de Cuba y Costa Rica. *Avicennia* **12/13**: 87–90.
- ORTEA, J. & ESPINOSA, J. 2001. Descripción de una nueva especie de *Ercolania* Trinchese, 1872. *Avicennia Suppl.* **4**: 45–47.
- ORTEA, J. & ESPINOSA, J. 2002. Nuevas especies del género *Elysia* Risso, 1818 (Mollusca: Sacoglossa) con caracteres singulares. *Avicennia* **15**: 129–140.
- ORTEA, J., LUQUE, A.A. & TEMPLADO, J. 1988. *Elysia picta* Verrill, 1901 and *Geitodoris pusae* (Marcus, 1955), two amphiatlantic opisthobranchs. *Journal of Molluscan Studies* **54**: 243–247.
- ORTEA, J. & MORO, L. 1998. Nota sobre *Ercolania siotti* Trinchese, 1872 (Mollusca: Opisthobranchia: Saccoglossa). *Revista de la Academia Canaria de Ciencias* **10**: 97–99.
- ORTEA, J., MORO, L., BACALLADO, J.J. & ESPINOSA, J. 1998. Catálogo abreviado de las especies del orden Sacoglossa (=Ascoglossa, Mollusca: Opisthobranchia) de las Islas Canarias y de Cabo Verde. *Revista de la Academia Canaria de Ciencias* **10**: 85–96.
- ORTEA, J., MORO, L. & ESPINOSA, J. 1997. Nuevos datos sobre el género *Elysia* Risso, 1818 (Opisthobranchia: Sacoglossa) en el Atlántico. *Revista de la Academia Canaria de Ciencias* **9**: 141–155.
- ORTEA, J. & TEMPLADO, J. 1988. Una nueva especie de *Cyerce* Bergh, 1871 (Opisthobranchia: Ascoglossa) de la Isla de Cuba. *Iberus* **8**: 11–14.
- ORTEA, J. & TEMPLADO, J. 1990. A new species of the genus *Cyerce* Bergh, 1871, from the Cape Verde Islands (Opisthobranchia: Ascoglossa). *The Veliger* **33**: 202–205.
- OSTERGAARD, J.M. 1955. Some opisthobranchiate Mollusca from Hawaii. *Pacific Science* **9**: 110–136.
- PEASE, W.H. 1861. Descriptions of new species of Mollusca from the Pacific Islands. *Proceedings of the Zoological Society, London*, **1861**: 242–247.
- PEASE, W.H. 1866. Remarks on Nudibranchiata inhabiting the Pacific islands, with descriptions of two new genera. *American Journal of Conchology* **2**: 204–208.
- PEASE, W.H. 1868. Descriptions of marine Gasteropoda inhabiting Polynesia. *American Journal of Conchology* **4**: 71–80, plates 7–12.
- PEASE, W.H. 1871. Descriptions of nudibranchiate Mollusca inhabiting Polynesia. *American Journal of Conchology* **6**: 299–305, plates 19–22.
- PERRONE, A.S. 1990. Una nuova specie di Elysiidae, *Elysia heta* nov. sp. dal litorale salentino (Mediterraneo, Golfo di Taranto) (Opisthobranchia: Sacoglossa). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano* **130**: 249–252.
- PIERCE, S.K., CURTIS, N.E., MASSEY, S.E., BASS, A.L., KARL, S.A. & FINNEY, C.M. 2006. A morphological and molecular comparison between *Elysia crispata* and a new species of kleptoplastic sacoglossan sea slug (Gastropoda: Opisthobranchia) from the Florida Keys, USA. *Molluscan Research* **26**: 23–38.
- PILSBRY, H.A. 1895. *Manual of conchology: structural and systematic*. **15**: 181–436, plates 43–50 & 59–61.
- PITTMAN, C. 2004. Re: *Bosellia* from Japan. *Message in Sea Slug Forum* (<http://www.seaslugforum.net/find.cfm?id=12348>) (last access 12th of August 2007).
- PLATTS, E. 1985. An annotated list of the North Atlantic Opisthobranchia. Appendix to: Just, H. and Edmunds, M. North Atlantic Nudibranchs (Mollusca) seen by Henning Lemeche. *Ophelia*, Supplement **2**: 150–170.
- POWELL, A.W.B. 1937. New species of nudibranchiate Mollusca from Auckland waters. *Records of the Auckland Institute Museum* **2**: 119–124.
- PRUD'HOMME VAN REINE, W.F., VERHEIJ, E. & COPPEJANS, E. 1996. Species and ecads of *Caulerpa* (Ulvophyceae, Chlorophyta) in Malesia (South-East Asia): Taxonomy, biogeography and biodiversity. *Netherlands Journal of Aquatic Ecology* **30**(2–3): 83–98.
- PRUVOT-FOL, A. 1953. Étude de quelques opisthobranches de la côte Atlantique du Maroc et du Senegal. *Travaux de l'Institut Scientifique Chérifien* **5**: 7–105, plates 1–3.
- PRUVOT-FOL, A. 1954. Mollusques opisthobranches. *Faune de France* **58**. Lechevalier, Paris.
- RAO, K.V. 1937. Structure, habits and early development of a new species of *Stiliger* Ehrenberg. *Records of the Indian Museum* **39**: 435–464, plates 7–9.
- RAO, K.V. & RAO, K.P. 1963. *Stiliger nigrovittatus* sp. nov., a sacoglossan mollusc from the Gulf of Mannar. *Journal of the Marine biological Association of India* **5**: 232–238.
- RIEK, D.W. 2006. First *Bosellia* from Australia. *Message in Sea Slug Forum* (<http://www.seaslugforum.net/find.cfm?id=17429>) (last access 12th of August 2007).
- RISBEC, J. 1928. Contribution à l'étude des nudibranches Néocalédoniens. *Faune de Colonies Françaises* **2**: 1–328.
- RISBEC, J. 1953. Mollusques nudibranches de la Nouvelle-Calédonie. *Faune de l'Union Française* **15**: 1–189.
- ROGINSKAYA, I. 2000. Russian Opisthobranchs. Notes on *Limapontia senestra* (Quatrefages, 1844) in White Sea and Barents Sea (Sacoglossa, Limapontiidae). *Nudibranch News* **2**(12): 56–58.
- RUDMAN, W.B. 2002. Comment on *Oxynoe viridis?* from Turkey by Baki Yokes. *Message in Sea Slug Forum* (<http://www.seaslugforum.net/find.cfm?id=6935>) (last access 12th of August 2007).
- SARMA, A.L.N. 1975. Three new species of the bivalved gastropods *Julia* and *Berthelinia* found in Eastern Indian Ocean. *Venus (Japanese Journal of Malacology)* **34**: 11–25.
- SCHMEKEL, L. & PORTMANN, A. 1982. *Opisthobranchia des Mittelmeeres. Nudibranchia und Saccoglossa*. Springer, Berlin - New York.
- SCHRÖDL, M. 1996. Nudibranchia y Sacoglossa de Chile: distribución, descripción externa y clave. *Gayana Zoologica* **60**: 17–62.
- SPENCER, H.G. & WILLAN, R.C. 1995. *The marine fauna of New Zealand: Index to the fauna 3. Mollusca*. New Zealand Oceanographic Institute, Wellington.
- SWENNEN, C. 1961. On a collection of Opisthobranchia from Turkey. *Zoologische Mededelingen* **38**: 41–75.
- SWENNEN, C. 1997. Two new gastropods, *Elysia bangtawensis* and *E. siamensis* from southern Thailand (Opisthobranchia, Sacoglossa, Elysiidae). *Bulletin Zoologisch Museum, Universiteit van Amsterdam* **16**(6): 33–39.
- SWENNEN, C. 2001. Two new sacoglossans (Gastropoda: Opisthobranchia) from Thailand. *Beaufortia Bulletin Zoological Museum University of Amsterdam* **51**(3): 75–81.
- SWENNEN, C., MOOLENBECK, R.G., RUTTANADAKUL, N., HOBBELINK, H., DEKKER, H. & HAJISAMAE, S. 2001. The Molluscs of the Southern Gulf of Thailand. *Thai Studies in Biodiversity No. 4*: 1–210.
- TEMPLADO, J., LUQUE, A.A. & ORTEA, J. 1990. A commented check-list of the amphiatlantic Ascoglossa and Nudibranchia (Mollusca: Opisthobranchia). *Lavori della Società Italiana di Malacologia, Napoli* **23**: 295–326.

- THOLLESSON, M. 1999. Phylogenetic analysis of Euthyneura (Gastropoda) by means of the 16S rRNA gene: the use of a 'fast' gene for 'higher-level' phylogenies. *Proceedings of the Royal Society of London, Series B* **266**: 75–83.
- THOMPSON, T.E. 1973. Sacoglossan gastropod molluscs from eastern Australia. *Proceedings of the Malacological Society of London* **40**: 239–251.
- THOMPSON, T.E. 1977. Jamaican opisthobranch molluscs I. *Journal of Molluscan Studies* **43**: 93–140.
- THOMPSON, T.E. 1979. Biology and relationships of the South African sacoglossan mollusc *Volvatella laguncula*. *Journal of Zoology, London* **189**: 339–47.
- THOMPSON, T.E. 1983. The Bermudan and Caribbean sacoglossan mollusk *Elysia flava* Verrill now recorded from the Greek Aegean Sea. *The Veliger* **26**: 136–138.
- THOMPSON, T.E. 1988. Eastern Mediterranean Opisthobranchia: Oxynoidae, Polybranchiidae, Stiligeridae (Sacoglossa). *Journal of Molluscan Studies* **54**: 157–172.
- THOMPSON, T.E. & JAKLIN, A. 1988. Eastern Mediterranean Opisthobranchia: Elysiidae (Sacoglossa = Ascoglossa). *Journal of Molluscan Studies* **54**: 59–69.
- THOMPSON, T.E., JARMAN, G.M. & ZENETOS, A. 1985. Infralittoral macrobenthos of the Patras Gulf and Ionian Sea: opisthobranch molluscs. *Journal of Conchology* **32**: 71–95.
- TROWBRIDGE, C.D. 1995a. Hypodermic insemination, oviposition, and embryonic development of a pool-dwelling ascoglossan (=sacoglossan) opisthobranch: *Ercolania felina* (Hutton, 1882) on New Zealand shores. *The Veliger* **38**: 203–211.
- TROWBRIDGE, C.D. 1995b. New Zealand opisthobranchs associated with the low intertidal, crustose green alga *Codium convolutum*: Ascoglossans "down under". *The Veliger* **38**: 116–125.
- TROWBRIDGE, C.D. 2002. Northeastern Pacific sacoglossan opisthobranchs: Natural history review, bibliography, and prospectus. *The Veliger* **45**(1): 1–24.
- TROWBRIDGE, C.D. 2006. Identity of *Stiliger akkeshiensis* from Kuril Islands. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=15995>) (last access 12th of August 2007).
- VADER, W. 1981. *Alderia modesta* (Gastropoda, Sacoglossa) in northern Norway. *Fauna Norvegica Seric A* **2**: 41–46.
- VALDÉS, A. & LOZOUET, P. 2000. Opisthobranch molluscs from the Tertiary of the Aquitaine Basin (south-western France), with descriptions of seven new species and a new genus. *Palaeontology* **43**(3): 457–479.
- VALDÉS, A., HAMANN, J., BEHRENS, D.W. & DUPONT, A. 2006. Caribbean Sea Slugs. A field guide to the opisthobranch mollusks from the tropical northwestern Atlantic. Sea Challengers Natural History Books, Etc., Gig Harbor, Washington, 289pp.
- VALENTINE, J.W. 1966. Numerical analysis of marine molluscan ranges on the extratropical northeastern Pacific shelf. *Limnology and Oceanography* **11**: 198–211.
- WÄGELE, H. & JOHNSON, G. 2001. Observations on the histology and photosynthetic performance of "solar-powered" opisthobranchs (Mollusca, Gastropoda, Opisthobranchia) containing symbiotic chloroplasts or zooxanthellae. *Organisms, Diversity and Evolution* **3**: 193–210.
- WÄGELE, H., VONNEMANN, V. & WÄGELE, J.W. 2003. Toward a phylogeny of the Opisthobranchia. Pp. 185–228 In: LYDEARD, C. & LINDBERG, D. (eds.) *Molecular Systematics and Phylogeography of Mollusks*. Smithsonian Institution Press, Washington, D.C.
- WAWRA, E. 1979. Zur systematischen Stellung von *Platyhedyle demudata* Salvini-Plawen, 1973 (Opisthobranchia, Gastropoda). *Zeitschrift für zoologische Systematik und Evolutionsforschung* **17**: 221–225.
- WILLAN, R. & MORTON, J. 1984. Marine Molluscs, part 2. Opisthobranchia. University of Auckland. Leigh Marine Laboratory: Leigh, 106pp.
- YOKES, B. 2001. *Elysia tomentosa* in the Mediterranean? Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=5676>) (last access 12th of August 2007).
- YOKES, B. 2002. *Elysia ornata?* from Turkey. Message in Sea Slug Forum (<http://www.seaslugforum.net/find.cfm?id=6932>) (last access 12th of August 2007).

Author's address: Zoological Museum, Universitetsparken 15, DK-2100 Copenhagen Ø, Denmark. E-mail: krjensen@snm.ku.dk.

APPENDIX

List of nominal species of recent Sacoglossa, authorship and type localities. Species are arranged alphabetically within families and superfamilies. Valid species (and species for which synonymy has been contested) are listed with their current generic name. Species considered invalid in the present study are marked with an *.

Species	Author	Type locality
Oxynoacea		
Volvatellidae		
<i>Ascobulla californica</i>	(Hamatani, 1971)	Gulf of California
<i>Ascobulla fischeri</i>	(Adams & Angas, 1864)	"South Australia"
<i>Ascobulla fragilis</i>	(Jeffreys, 1856)	Mediterranean
<i>Ascobulla japonica</i>	(Hamatani, 1969)	Kii, Middle Japan
<i>Ascobulla? pusilla</i>	(Nevill & Nevill, 1869)	Sri Lanka
<i>Ascobulla souverbiei</i>	(Montrouzier in Souverbie & Montrouzier, 1874)	New Caledonia
<i>Ascobulla? systemna*</i>	(Melvill, 1918)	Gulf of Oman
<i>Ascobulla ulla</i>	(Marcus & Marcus, 1970)	E of Santos, Brazil
<i>Ascobulla? turtoni*</i>	(Bartsch, 1915)	South Africa
<i>Volvatella angeliniana</i>	Ichikawa, 1993	Sesoko Isl., Ryukyu
<i>Volvatella australis</i>	Jensen, 1997	Darwin Harbour, N Australia
<i>Volvatella ayakii</i>	Hamatani, 1972	Kii, Middle Japan
<i>Volvatella bermudae</i>	Clark, 1982	Bermuda
<i>Volvatella candida</i>	Pease, 1868	French Polynesia
<i>Volvatella cincta</i>	Nevill & Nevill, 1869	Sri Lanka
<i>Volvatella elioti</i>	(Evans, 1950)	Zanzibar
<i>Volvatella evansi</i>	(Kay, 1961)	Oahu, Hawaii
<i>Volvatella ficula</i>	Burn, 1966	Fiji
<i>Volvatella fragilis</i>	Pease, 1860	Sandwich Islands
<i>Volvatella kawamurai*</i>	Habe, 1946	Japan
<i>Volvatella laguncula</i>	Sowerby, 1894	Port Elizabeth, S Africa
<i>Volvatella omega</i>	(Melvill, 1918)	Gulf of Oman
<i>Volvatella pyriformis</i>	Pease, 1868	Huaheine, French Polynesia
<i>Volvatella ventricosa</i>	Jensen & Wells, 1990	Albany, SW Australia
<i>Volvatella vigourouxi</i>	(Montrouzier, 1861)	New Caledonia
<i>Volvatella viridis</i>	Hamatani, 1976	Anami Islands, Japan
Juliidae		
<i>Berthelinia (Midorigai) australis</i>	(Burn, 1960)	Torquay, Victoria, Australia
<i>Berthelinia (Tamanovalva) babai</i>	(Burn, 1965)	Torquay, Victoria, Australia
<i>Berthelinia caribbea</i>	Edmunds, 1963	Port Royal, Jamaica
<i>Berthelinia chloris</i>	(Dall, 1918)	Baja California, W Mexico
<i>Berthelinia corallensis*</i>	Hedley, 1920	Queensland, Australia
<i>Berthelinia darwini</i>	Jensen, 1997	Darwin Harbour, N Australia
<i>Berthelinia (Tamanovalva) fijiensis</i>	(Burn, 1966)	Fiji
<i>Berthelinia ganapati</i>	Sarma, 1975	SE India
<i>Berthelinia (Tamanovalva) limax</i>	(Kawaguti & Baba, 1959)	Seto, Japan
<i>Berthelinia pseudochloris</i>	Kay, 1964	Kauai, Hawaii
<i>Berthelinia rotnesti</i>	Jensen, 1993	Rottnest Island, SW Australia
<i>Berthelinia schlumbergeri</i>	Dautzenberg, 1895	Madagascar
<i>Berthelinia (Edentellina) typica</i>	(Gatliff & Gabriel, 1911)	Port Phillip, Victoria, Australia
<i>Berthelinia waltairensis</i>	Sarma, 1975	SE India
<i>Julia borbonica*</i>	(Deshayes, 1863)	Reunion
<i>Julia burni</i>	Sarma, 1975	Andaman Islands, India
<i>Julia cornuta*</i>	(De Folin, 1867)	Mauritius
<i>Julia equatorialis*</i>	Pilsbry & Olsson, 1944	N of Mancara, Peru
<i>Julia exquisita</i>	Gould, 1862	Hawaiian Islands
<i>Julia japonica</i>	Kuroda & Habe, 1951	Wakayama, Honshu, Japan
<i>Julia mishimaensis</i>	Kawaguti & Yamasu, 1982	Yamaguchi Pref. Japan
<i>Julia thecaphora</i>	(Carpenter, 1857)	Mazatlán, Mexico(?)
<i>Julia zebra</i>	Kawaguti 1981	Yamaguchi Pref. Japan
Oxynoidea		
<i>Icarus gravesii*</i>	Forbes, 1844	Aegcan Sea
<i>Lobiger corneus*</i>	Mörch, 1863	? (Cuming collection)

<i>Lobiger cumingi</i> *	(A. Adams, 1850)	Puerto St. Elena, W Colombia
<i>Lobiger uevilli</i>	Pilsbry, 1896	n.n. for <i>L. viridis</i> G. & H. Nevill, 1869
<i>Lobiger pellicidus</i> *	A. Adams, 1854	unknown (Cuming collection)
<i>Lobiger philippi</i> *	Krohn, 1847	Sicily, Italy
<i>Lobiger pilsbryi</i> *	Schwengel, 1941	Sanibel Isl., Florida
<i>Lobiger picta</i> *	Pease, 1868	Huaheine, French Polynesia
<i>Lobiger sagamiensis</i>	Baba, 1952	Sagami Bay, Japan
<i>Lobiger serradifalci</i>	(Calcara, 1840)	Sicily, Italy
<i>Lobiger souverbii</i>	Fischer, 1856	Guadeloupe
<i>Lobiger viridis</i>	Pease, 1863	Huahine
<i>Lobiger viridis</i> *	Nevill & Nevill, 1869	Sri Lanka
<i>Lophocercus krolmii</i> *	A. Adams, 1854	Hawaiian Islands (Sandwich Islands)
<i>Lophocercus sieboldii</i> *	Krohn, 1847	Messina, Italy
<i>Lophopleurella capensis</i>	(Thiele, 1912)	S Africa
<i>Oxynoe aguayoi</i> *	Jaume, 1945	Cuba
<i>Oxynoe antillarum</i>	Mörch, 1863	St. Thomas, USVI
<i>Oxynoe azuroopunctata</i>	Jensen, 1980	Florida
<i>Oxynoe benchijigua</i>	Ortea, Moro & Espinosa, 1999	Canary Islands
<i>Oxynoe brachycephalus</i> *	Mörch, 1863	Based on figure
<i>Oxynoe delicatula</i>	Nevill & Nevill, 1869	Sri Lanka
<i>Oxynoe hargravesi</i> *	Adams, 1872	New Hebrides
<i>Oxynoe kabirensis</i>	Hamatani, 1980	Ishigaki Isl., Ryukyu Islands
<i>Oxynoe natalensis</i> *	Smith, 1903	Port Elizabeth, S Africa
<i>Oxynoe olivacea</i>	Rafinesque, 1814	Sicily, Italy
<i>Oxynoe panamensis</i>	Pilsbry & Olsson, 1943	Bocas Isl., Panama
<i>Oxynoe viridis</i>	(Pease, 1861)	Sandwich Islands
<i>Roburnella wilsoni</i>	(Tate, 1889)	Port Phillip Bay, Victoria, Australia
Plakobranchacea		
Plakobranchidae		
<i>Actaeon elegans</i> *	de Quatrefages, 1844	St. Vaast, France
<i>Aplysiopterus neapolitanus</i> *	delle Chiaje, 1830	Napoli, Italy
<i>Elisia marmorata</i> *	Cantraine, 1835	Livorno, Italy
<i>Elisia abei</i>	Baba, 1955	Sagami Bay, Japan
<i>Elisia albomarginata</i> *	Trinchese, 1869	Italy
<i>Elisia amakusana</i>	Baba, 1955	Sagami Bay, Japan
<i>Elisia atroviridis</i>	Baba, 1955	Sagami Bay, Japan
<i>Elisia australis</i>	(Quoy & Gaimard, 1832)	Port Jackson, Sydney, NSW
<i>Elisia babai</i>	Pruvot-Fol, 1946	Ryukyu Islands
<i>Elisia bangtawanaensis</i>	Swennen, 1997	Pattani, Gulf of Thailand
<i>Elisia bedecta</i> *	MacFarland, 1966	Monterey Bay, California
<i>Elisia bella</i> ³	(Pease, 1860)	Hawaii
<i>Elisia bennettiae</i>	Thompson, 1973	Heron Island, GBR
<i>Elisia canguzua</i>	Marcus, 1955	NE of Santos, Brazil
<i>Elisia catulus</i>	(Gould, 1870)	Massachusetts, USA
<i>Elisia cauze</i> *	Marcus, 1957	NE of Santos, Brazil
<i>Elisia chilkenis</i>	Eliot, 1916	Chilka Lake, India
<i>Elisia chitwa</i>	Marcus, 1955	NE of Santos, Brazil
<i>Elisia chlorotica</i>	Gould, 1870	Massachusetts, USA
<i>Elisia clarki</i>	Pierce et al., 2006	Florida
<i>Elisia clena</i> *	Marcus & Marcus, 1970	?Curacao/?Barbados
<i>Elisia coodgeensis</i>	(Angas, 1864)	Port Jackson, Sydney, NSW
<i>Elisia cornigera</i> *	Nuttall, 1989	Spanish Harbor Key, FL
<i>Elisia crispata</i>	Mörch, 1863	St. Croix, USVI
<i>Elisia cyanea</i> *	Mamo in Caruana, 1867	Malta
<i>Elisia degeneri</i>	Ostergaard, 1955	Oahu, Hawaii
<i>Elisia diomedea</i>	(Bergh, 1894)	Lower California
<i>Elisia dubia</i> *	Eliot, 1904	Zanzibar
<i>Elisia duiis</i> *	Marcus & Marcus, 1967	Biscayne Bay, Florida
<i>Elisia elsiae</i> *	Ostergaard, 1955	Waikiki, Hawaii
<i>Elisia eugeniae</i>	Ortea & Espinosa, 2002	Manzanillo, Costa Rica (Carib)
<i>Elisia evelinae</i>	Marcus, 1957	NE of Santos, Brazil
<i>Elisia kushimotoensis</i>	Baba, 1957	Kushimoto, Kii, Japan
<i>Elisia expansa</i>	(O'Donoghue, 1924)	Abrolhos Islands, NW Australia
<i>Elisia faustula</i> *	Bergh, 1872	Masoloc, Philippines

<i>Elysia fezi</i>	Vilella, 1968	Barcelona, Spain
<i>Elysia filicauda</i>	Jensen & Wells, 1990	Albany, SW Australia
<i>Elysia flava</i>	Verrill, 1901	Bermuda
<i>Elysia flavipunctata</i>	Ichikawa, 1993	Ishigaki Isl., Ryukyu
<i>Elysia flavomacnla</i>	Jensen, 1990	Hong Kong
<i>Elysia furvacauda</i>	Burn, 1958	Torquay, Victoria, Australia
<i>Elysia fusca*</i>	Philippi, 1844	Palermo, Italy
<i>Elysia gordanae</i>	Thompson & Jaklin, 1988	Rovinj, Yugoslavia
<i>Elysia grandifolia</i>	Kelaart, 1857	Sri Lanka
<i>Elysia grandis³</i>	Bergh, 1872	Palau? (Pclew)
<i>Elysia haingsisiana³</i>	Bergh, 1905	Haingsisi
<i>Elysia halimeda*</i>	Macnae, 1954	Port Alfred, S Africa
<i>Elysia hamatani</i>	Baba, 1957	Seto, Japan
<i>Elysia hedgpethi</i>	Marcus, 1961	Tomales Bay, California
<i>Elysia hendersoni</i>	Eliot, 1899	Samoa
<i>Elysia hetta</i>	Perrone, 1990	Gulf of Taranto, Italy
<i>Elysia hirasei</i>	Baba, 1955	Sagami Bay, Japan
<i>Elysia japonica</i>	Eliot, 1913	Japan
<i>Elysia latipes*</i>	Marcus, Er. & Marcus, Ev. 1960	Maldives
<i>Elysia lencolegnote</i>	Jensen, 1990	Hong Kong
<i>Elysia lobata</i>	Gould, 1852	Hawaii
<i>Elysia macnaei*</i>	Ev. Marcus, 1980	Japan?/Australia?
<i>Elysia maoria</i>	Powell, 1937	Auckland, NZ
<i>Elysia margaritae</i>	Fez, 1962	Valencia, Spain
<i>Elysia marginata*</i>	(Pease, 1871)	Huachine, French Polynesia
<i>Elysia minima</i>	Ichikawa, 1993	Kuro Isl., Ryukyu
<i>Elysia mimia*</i>	(Sars, 1835)	Bergensund, Norway
<i>Elysia nealae</i>	Ostergaard, 1955	Waikiki, Hawaii
<i>Elysia nigrocapitata</i>	Baba, 1957	?Osaka/?Seto/?Tsurugu/?Toyama Bay
		Japan
<i>Elysia nigropunctata</i>	(Pease, 1871)	Tahiti
<i>Elysia nisbeti</i>	Thompson, 1977	Jamaica
<i>Elysia obtusa</i>	Baba, 1938	Seto, Japan
<i>Elysia ocellata*</i>	Pease, 1860	Hawaii
<i>Elysia oerstedii</i>	Morch, 1859	Puntarenas, Central America
<i>Elysia ornata</i>	(Swainson, 1840)	West Indies
<i>Elysia ornata*</i>	(Pease, 1860)	Hawaii
<i>Elysia pagenstecheri*</i>	Marcus, Ev., 1982	Sète, Mediterranean
<i>Elysia papillosa</i>	Verrill, 1901	Bermuda
<i>Elysia patagonica</i>	Munian & Ortea, 1997	San Jorge Gulf, Argentina (45d58'S; 67d34'W)
<i>Elysia patina</i>	Marcus, 1980	Florida Keys
<i>Elysia pilosa</i>	Risbec, 1928	New Caledonia
<i>Elysia pratensis</i>	Ortea & Espinosa, 1996	Eastern part of Yucatan, Mexico
<i>Elysia pruvotae*</i>	Risbec, 1953	New Caledonia
<i>Elysia punctata</i>	Kelaart, 1857	Sri Lanka
<i>Elysia purchoni</i>	Thompson, 1977	Jamaica
<i>Elysia rufescens</i>	(Pease, 1871)	Tahiti
<i>Elysia serca</i>	Marcus, 1955	NE of Santos, Brazil
<i>Elysia setoensis</i>	Hamatani, 1968	Seto, Japan
<i>Elysia siamensis</i>	Swennen, 1997	Pattani, Gulf of Thailand
<i>Elysia slimora</i>	Marcus & Marcus, 1966	near Sao Tome, W Africa
<i>Elysia splendida*</i>	Grube, 1861	Cherso, N Adriatic
<i>Elysia subornata</i>	Verrill, 1901	Bermuda
<i>Elysia sugashimae</i>	Baba, 1955	Sagami Bay, Japan
<i>Elysia thompsoni</i>	Jensen, 1993	Rottne Island, SW Australia
<i>Elysia timida</i>	Risso, 1818	Nice, France Mediter.
<i>Elysia tokarensis</i>	Baba, 1957	Tokara Islands, Kyushu, Japan
<i>Elysia tomentosa</i>	Jensen, 1997	Abrolhos Islands, NW Australia
<i>Elysia translucens</i>	Pruvot-Fol, 1957	Banyuls, France Mediter.
<i>Elysia trilobata</i>	Heller & Thompson, 1983	Red Sea
<i>Elysia trisimata</i>	Baba, 1949	Sagami Bay, Japan
<i>Elysia tuca</i>	Marcus & Marcus, 1967	Biscayne Bay, Florida
<i>Elysia (Elysiella) verrilli*</i>	Thiele, 1931	n.n. for <i>Elysia (Elysiella) catula</i> Verrill
<i>Elysia verrucosa</i>	Jensen, 1985	Hong Kong

<i>Elysia viridis</i>	(Montagu, 1804)	Devonshire, UK
<i>Elysia viridissima</i> *	Trinchese, 1869	Italy
<i>Elysia vreelandae</i>	Marcus & Marcus, 1970	W Mexico
<i>Elysia yocoyamana</i>	Baba, 1936	Ishigakishima, Ryukyu
<i>Elysia zuleicae</i>	Ortea & Espinosa, 2002	Cuba
<i>Elysiella pusilla</i>	Bergh, 1872	Aibukit, ?Palau (Palaos)
<i>Elysiella stylifera</i>	Jensen, 1997	Darwin Harbour, N Australia
<i>Elysiobranchus mercieri</i>	Pruvot-Fol, 1930	New Caledonia
<i>Elysiobranchus ryukyuensis</i>	Ichikawa, 1993	Sesoko Isl., Ryukyu
<i>Pattyclaya arena</i>	Carlson & Hoff, 1978	Guam
<i>Pattyclaya brycei</i>	Jensen & Wells, 1990	Albany, SW Australia
<i>Placobranchus argus</i> *	Bergh, 1872	Honolulu, Hawaii
<i>Placobranchus caminguinus</i> *	Bergh, 1872	Luzon, Philippines
<i>Placobranchus chlorophagus</i> *	Bergh, 1878	Huaheine
<i>Placobranchus gracilis</i> *	Pease, 1871	Tahiti
<i>Placobranchus guttatus</i> *	Stimpson, 1858	Ryukyu Islands
<i>Plakobranchus ianthobapsus</i> *	Gould, 1852	Honolulu, Hawaii
<i>Plakobranchus laetus</i> *	Bergh, 1872	Masoloc, Philippines
<i>Plakobranchus ocellatus</i>	van Hasselt, 1824	Sunda Strait
<i>Plakobranchus priapinus</i> *	Bergh, 1872	Bohol, Philippines
<i>Plakobranchus punctulatus</i> *	Bergh, 1872	Masoloc, Philippines
<i>Plakobranchus variegatus</i> *	Pease, 1871	Huaheine, French Polynesia
<i>Thuridilla albopustulosa</i>	Gosliner, 1995	Madang, PNG
<i>Thuridilla bayeri</i>	(Marcus, 1965)	Marshall Islands
<i>Thuridilla carlsoni</i>	Gosliner, 1995	Madang, PNG
<i>Thuridilla coerulea</i>	(Kelaart, 1857)	Sri Lanka
<i>Thuridilla decorata</i>	(Heller & Thompson, 1983)	Red Sea
<i>Thuridilla flavomaculata</i>	Gosliner, 1995	Luzon, Philippines
<i>Thuridilla gracilis</i>	(Risbee, 1928)	New Caledonia
<i>Thuridilla hoffae</i>	Gosliner, 1995	Madang, PNG
<i>Thuridilla hopei</i>	(Verany, 1853)	Nice, France Mediterr.
<i>Thuridilla indopacifica</i>	Gosliner, 1995	Aldabra Atoll
<i>Thuridilla kathae</i>	Gosliner, 1995	Madagascar
<i>Thuridilla lineolata</i>	(Bergh, 1905)	Saleyer
<i>Thuridilla livida</i>	(Baba, 1955)	Sagami Bay, Japan
<i>Thuridilla mazda</i>	Ortea & Espinosa, 2000	Manzanillo, Costa Rica (Carib)
<i>Thuridilla moebii</i>	(Bergh, 1888)	Mauritius
<i>Thuridilla multimarginata</i>	Gosliner, 1995	Midway Atoll
<i>Thuridilla neona</i>	Gosliner, 1995	Lanai, Hawaii
<i>Thuridilla picta</i>	(Verrill, 1901)	Bermuda
<i>Thuridilla ratna</i>	(Marcus, 1965)	Palau
<i>Thuridilla splendens</i>	(Baba, 1949)	Sagami Bay, Japan
<i>Thuridilla thysanopoda</i> ³	(Bergh, 1905)	Tual, Kei Islands, Indonesia
<i>Thuridilla undula</i>	Gosliner, 1995	Madang, PNG
<i>Thuridilla vatiae</i>	(Risbee, 1928)	New Caledonia
<i>Thuridilla virgata</i>	(Bergh, 1888)	Mauritius
<i>Tridachia schrammi</i> *	Mörch, 1863	Guadeloupe
Boselliidae		
<i>Bosellia cohellia</i> ²	Marcus, 1978	?Red Sea/?Mediterranean
<i>Bosellia coriunae</i>	Marcus, 1973	Key Biscayne, Florida
<i>Bosellia leve</i>	Fernandez-Ovies & Ortea, 1986	Lanzarote, Canary Islands
<i>Bosellia marcusii</i>	Marcus, 1972	Grassy Key & Miami, FL
<i>Bosellia mimetica</i>	Trinchese, 1890	Capri, Italy
Platyhedylidae		
<i>Gascoignella aprica</i>	Jensen, 1985	Hong Kong
<i>Gascoignella jabae</i>	Swennen, 2001	Pattani, Gulf of Thailand
<i>Gascoignella mukuli</i>	Swennen, 2001	Pattani, Gulf of Thailand
<i>Platyhedyle denudata</i>	Salvini-Plawen, 1973	Livorno, Italy
Limapontioidea		
Polybranchidae (=Caliphyllidae)		
<i>Caliphylla mediterranea</i>	(A. Costa, 1867)	Napoli, Italy
<i>Caliphylla tricolor</i> *	Trinchese, 1879	Mediterranean
<i>Cyerce antillensis</i>	Engel, 1927	Tobago(?), Westindien

<i>Cyerce cristallina</i>	(Trinchese, 1881)	Napoli, Italy
<i>Cyerce? edmundsi</i>	Thompson, 1977	Jamaica
<i>Cyerce elegans</i>	Bergh, 1871	Palau? (Palaos)
<i>Cyerce graeca</i>	Thompson, 1988	Greek Ionian Sea
<i>Cyerce habanensis</i>	Ortea & Templado, 1988	Cuba
<i>Cyerce jheringi*</i>	Pelsneer, 1892	Napoli, Italy
<i>Cyerce kikutarobabai</i>	Hamatani, 1976	Amami Islands, Japan
<i>Cyerce nigra</i>	Bergh, 1871	Palau? (Palaos)
<i>Cyerce nigricans</i>	(Pease, 1866)	Pacific Islands
<i>Cyerce orzeai</i>	Valdès & Camacho, 2000)	Puntarenas, Costa Rica
<i>Cyerce pavonina</i>	Bergh, 1888	Mauritius
<i>Cyerce verdensis</i>	Ortea & Templado 1990	Capc Verde
<i>Mourgona germaineae</i>	Marcus & Marcus, 1970	Puerto Rico (aquarium)
<i>Mourgona murca</i>	Marcus & Marcus, 1970	Curacao
<i>Mourgona osuni</i>	Hamatani, 1994	Amami-Oshima Island, SW Japan
<i>Polybranchia borgnini</i>	(Trinchese, 1895)	Mediterranean
<i>Polybranchia orientalis</i>	(Kelaart, 1858)	Sri Lanka
<i>Polybranchia pallens</i>	(Burn, 1957)	Queenscliff, Victoria, Australia
<i>Polybranchia papillosa</i>	(Pease, 1866)	Pacific Islands
<i>Polybranchia pellucida</i>	Pease, 1860	Hawaii
<i>Polybranchia prasinus</i>	(Bergh, 1871)	Luzon, Philippines
<i>Polybranchia rubicundus*</i>	(Bergh, 1871)	Tor, Red Sea
<i>Polybranchia viridis</i>	(Deshayes, 1857)	Guadeloupe
<i>Polybranchia westralis</i>	Jensen, 1993	Rottneest Island, SW Australia
<i>Sohgenia palauensis</i>	Hamatani, 1991	Palau
Hermaeidae		
<i>Aplysiopsis brattstroemi</i>	(Marcus, 1959)	Chile (23d39'S; 70d25'W)
<i>Aplysiopsis elegans</i>	(Deshayes, 1835)	Banyuls, France Mediterr.
<i>Aplysiopsis enteromorphae</i>	(Cockerell & Eliot, 1905)	San Pedro, California(?)
<i>Aplysiopsis formosa</i>	Pruvot-Fol, 1953	Temara, Morocco
<i>Aplysiopsis maculosa*</i>	(Trinchese, 1874)	Genova, Italy
<i>Aplysiopsis minor</i>	(Baba, 1959)	Toyama Bay, Japan
<i>Aplysiopsis nigra</i>	(Baba, 1949)	Sagami Bay, Japan
<i>Aplysiopsis orientalis</i>	(Baba, 1949)	Sagami Bay, Japan
<i>Aplysiopsis sinusmensalis</i>	(Macnae, 1954)	Table Bay, S Africa
<i>Aplysiopsis smithi*</i>	(Marcus, 1961)	Tomales Bay, California
<i>Aplysiopsis toyamana</i>	(Baba, 1959)	Toyama Bay, Japan
<i>Aplysiopsis zebra*</i>	Clark, 1982	Florida
<i>Hermaea bifida</i>	(Montagu, 1816)	Devonshire, UK
<i>Hermaea boucheti</i>	Cervera, Garcia-Gomez & Ortea, 1991	Pontevedra, SW Spain
<i>Hermaea carminis*</i>	Fez, 1962	Valencia, Spain
<i>Hermaea coirala</i>	Marcus, 1955	NE of Santos, Brazil
<i>Hermaea cruciata</i>	Gould, 1870	Massachusetts, USA
<i>Hermaea evelinemarcusae</i>	Jensen, 1993	Rottneest Island, SW Australia
<i>Hermaea hillae</i>	Marcus & Marcus, 1967	Sonora, W Mexico
<i>Hermaea lutescens*</i>	A. Costa, 1866	Napoli, Italy
<i>Hermaea minor*</i>	Bergh, 1888	Mauritius
<i>Hermaea noto</i>	(Baba, 1959)	?Noto Peninsula/?Toyama Bay Japan
<i>Hermaea oliviae</i>	(MacFarland, 1966)	Monterey Bay, California
<i>Hermaea paucicirra</i>	Pruvot-Fol, 1953	Marocco
<i>Hermaea polychroma*</i>	(Hesse, 1873)	Brest
<i>Hermaea vancouverensis</i>	O'Donoghue, 1924	Vancouver Isl., Canada
<i>Hermaea variopicta</i>	(A. Costa, 1869)	Napoli, Italy
<i>Hermaea venosa*</i>	Lovén, 1845	Bohuslän, Sweden
<i>Hermaea wrangeliae</i>	(Ichikawa, 1993)	Kuro Isl., Ryukyu
<i>Hermaea zosteriae</i>	(Baba, 1959)	Amakusa, Japan
<i>Physopneumon carneum*</i>	A. Costa, 1862	Mediterranean
Limapontiidae		
<i>Alderella comosa</i>	(Costa, 1867)	Napoli, Italy
<i>Alderella harvardiensis*</i>	Gould, 1870	Massachusetts, USA
<i>Alderella modesta</i>	(Lovén, 1844)	Bohuslän, Sweden
<i>Alderella scaldiana*</i>	Nyst, 1855	Scheldt estuary, Netherlands
<i>Alderella uda</i>	Marcus, Ev. & Marcus, Er., 1956)	SW of Santos, Brazil
<i>Alderella willowi</i>	Krug, Ellingson, Burton & Valdés, 2007	San Pedro, California

<i>Alderiopsis garfio</i>	Caballer, Ortea & Espinosa, 2006	Cuba
<i>Alderiopsis nigra</i>	(Baba, 1937)	Amakusa, Japan
<i>Calliopaea bellula</i>	d'Orbigny, 1837	France
<i>Calliopaea oophaga</i>	Lemche, 1974	Samsø, Kattegat, DK
<i>Calliopaea souleyeti*</i>	Verany, 1853	Nice, France Mediterr.
<i>Cenia cocksii*</i>	Alder & Hancock, 1848	Falmouth, UK
<i>Cenia corrugata*</i>	Alder & Hancock, 1848	Falmouth, UK
<i>Chalidis coerules*</i>	de Quatrefages, 1844	Ile de Bréhat, France
<i>Chalidis nigricans*</i>	Alder & Hancock, 1847	Falmouth, UK
<i>Costasiella formicaria</i>	(Baba, 1959)	Amakusa, Japan
<i>Costasiella illa</i>	(Marcus, 1965)	Caroline Islands, Micronesica
<i>Costasiella iridophora</i>	Ichikawa, 1993	Kuro Isl., Ryukyu
<i>Costasiella kuroshimae</i>	Ichikawa, 1993	Kuro Isl., Ryukyu
<i>Costasiella lilianae*</i>	(Marcus, Ev. & Marcus, Er., 1969)	NE of Santos, Brazil
<i>Costasiella mandurahae</i>	Jensen, 1997	Darwin Harbour, N Australia
<i>Costasiella nonatoi</i>	Marcus & Marcus, 1960	NE of Santos, Brazil
<i>Costasiella ocellifera</i>	(Simroth, 1895)	Bermuda
<i>Costasiella pallida</i>	Jensen, 1985	Hong Kong
<i>Costasiella paweli</i>	Ichikawa, 1993	Miyako Isl., Ryukyu
<i>Costasiella rubrolineata</i>	Ichikawa, 1993	Ishigaki Isl., Ryukyu
<i>Costasiella usagi</i>	Ichikawa, 1993	Ishigaki Isl., Ryukyu
<i>Costasiella vegae</i>	Ichikawa, 1993	Ishigaki Isl., Ryukyu
<i>Costasiella virescens</i>	Pruvot-Fol, 1951	Banyuls, France Mediterr.
<i>Custiphorus vesiculosus*</i>	Deshaycs, 1853	?
<i>Embletonia mariae*</i>	Meyer & Möbius, 1865	Kieler Bucht
<i>Ercolania boodleae</i>	(Baba, 1938)	Seto, Japan
<i>Ercolania coerulea</i>	Trinchese, 1892	Napoli, Italy
<i>Ercolania costai*</i>	Pruvot-Fol, 1951	?Banyuls/?Monaco, Mediterranean
<i>Ercolania cricetae*</i>	(Marcus & Marcus, 1970)	Curacao
<i>Ercolania evelinae</i>	(Marcus, 1959)	Chile (53d22'S; 70d57'W)
<i>Ercolania emarginata*</i>	Jensen, 1985	Hong Kong
<i>Ercolania endophytophaga</i>	Jensen, 1999	Rottneest Island, SW Australia
<i>Ercolania erbsus</i>	(Marcus & Marcus, 1970)	Madagascar
<i>Ercolania felina</i>	(Hutton, 1882)	New Zealand
<i>Ercolania funerea*</i>	(Costa, 1867)	Napoli, Italy
<i>Ercolania fuscata</i>	(Gould, 1870)	Massachusetts, USA
<i>Ercolania gopalai</i>	(Rao, 1937)	Madras, E India
<i>Ercolania irregularis</i>	(Eliot, 1904)	Zanzibar
<i>Ercolania lozanoi</i>	Ortea, 1981	Tenerife, Canary Islands
<i>Ercolania margaritae</i>	Burn, 1974	Point Lonsdale, Victoria, Australia
<i>Ercolania nigra</i>	(Lemche, 1936)	Nyborg Fjord, DK
<i>Ercolania nigrovittata*</i>	(A. Costa, 1866)	Napoli, Italy
<i>Ercolania nigrovittata*</i>	(Rao & Rao, 1963)	Gulf of Mannar, SE India
<i>Ercolania pancerii*</i>	Trinchese, 1872	Genova, Italy
<i>Ercolania pica</i>	(Annandale & Prashad, 1922)	Chilka Lake, India
<i>Ercolania raorum</i>	(Marcus & Marcus, 1970)	Gulf of Mannar, SE India
<i>Ercolania selva</i>	Ortea & Espinosa, 2001	Manzanillo, Costa Rica (Carib)
<i>Ercolania siottii</i>	Trinchese, 1872	Genova, Italy
<i>Ercolania subviridis</i>	(Baba, 1959)	Toyama Bay, Japan
<i>Ercolania talis*</i>	(Marcus & Marcus, 1956)	SW of Santos, Brazil
<i>Ercolania tentaculata</i>	(Eliot, 1917)	Singgora, SE Thailand
<i>Ercolania translucens</i>	Jensen, 1993	Rottneest Island, SW Australia
<i>Ercolania trincheseii*</i>	Pruvot-Fol, 1951	?Banyuls/?Monaco, Mediterranean
<i>Ercolania uziellii*</i>	Trinchese, 1872	Genova, Italy
<i>Ercolania vanellus*</i>	Marcus, 1957	E of Santos, Brazil
<i>Ercolania varians</i>	(Eliot, 1904)	Zanzibar
<i>Ercolania viridis</i>	(A. Costa, 1866)	Napoli, Italy
<i>Ercolania zanzibarica</i>	Eliot, 1903	E Zanzibar
<i>Limapontia capitata</i>	(Müller, 1773)	Baltic Sea
<i>Limapontia cornuta*</i>	Giard, 1873	?
<i>Limapontia depressa</i>	Alder & Hancock, 1862	Sunderland, UK
<i>Limapontia nigra*</i>	Johnston, 1836	Berwick Bay, UK
<i>Limapontia senestra</i>	(de Quatrefages, 1844)	Ile de Bréhat, France
<i>Limapontia zonata</i> ¹	(Girard, 1852)	Massachusetts, USA
<i>Olea hansineensis</i>	Agersborg, 1923	Friday Harbor, Washington

<i>Placida aoteana</i> *	(Powell, 1937)	Auckland, NZ
<i>Placida babai</i> *	Mareus, Ev., 1982	Seto, Japan
<i>Placida brevicornis</i>	(A. Costa, 1867)	Napoli, Italy
<i>Placida capensis</i> *	Macnac, 1954	Cape Province, S Africa
<i>Placida cremoniana</i>	(Trinchese, 1892)	Napoli, Italy
<i>Placida daguilarensis</i>	Jensen, 1990	Hong Kong
<i>Placida dakariensis</i>	(Pruvot-Fol, 1953)	Dakar, Senegal
<i>Placida dendritica</i>	(Alder & Hancock, 1843)	Torbay, UK
<i>Placida fralila</i>	Burn, 1966	Queensland, Australia
<i>Placida kingstoni</i>	Thompson, 1977	Jamaica
<i>Placida ornata</i> *	(MacFarland, 1966)	Monterey Bay, California
<i>Placida saronica</i>	(Thompson, 1988)	Greek Aegean Sea
<i>Placida tardyi</i>	(Trinchese, 1873)	Genova, Italy
<i>Placida verticillata</i>	Ortea, 1981	Tenerife, Canary Islands
<i>Placida viridis</i>	(Trinchese, 1873)	Genova, Italy
<i>Stiliger akkeshiensis</i>	Baba, 1935	Akkeshi Bay, Hokkaido, Japan
<i>Stiliger aureomarginatus</i>	Jensen, 1993	Rottneest Island, SW Australia
<i>Stiliger bergli</i>	Baba, 1937	Tomioka, Amakusa, Japan
<i>Stiliger fuscovittatus</i>	Lancee, 1962	San Diego, California
<i>Stiliger llerai</i>	Ortea, 1981	Tenerife, Canary Islands
<i>Stiliger pusillus</i>	Baba, 1959	Osaka Bay, Japan
<i>Stiliger smaragdinus</i>	Baba, 1949	Sagami Bay, Japan
<i>Stiliger ornatus</i>	Ehrenberg, 1828	Red Sea
<i>Stiliger vossi</i>	Mareus & Marcus, 1960	Upper Florida Keys
<i>Stiliger? viridis</i> ²	(Kelaart, 1858)	Sri Lanka

¹This species is probably a flatworm rather than a sacoglossan.

²The taxonomic status of this species is so uncertain that it has been omitted from the analyses.

³This species has been omitted from the analyses due to lack of information.

⁴Since the type locality is uncertain, this species has been omitted from the analyses.

ZOBODAT - www.zobodat.at

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Bonn zoological Bulletin - früher Bonner Zoologische Beiträge.](#)

Jahr/Year: 2007

Band/Volume: [55](#)

Autor(en)/Author(s): Jensen Kathe R.

Artikel/Article: [Biogeography of the Sacoglossa \(Mollusca, Opisthobranchia\) 255-281](#)