

## **Vegetation of screes of the montane and colline zones in the Pamir-Alai Mts in Tajikistan (Middle Asia)**

### **Die Schuttvegetation der montanen und kollinen Zone des Pamir-Alai-Gebirges in Tadschikistan (Mittelasien)**

Arkadiusz Nowak<sup>1,2,\*</sup>, Sylwia Nowak<sup>1</sup>, Marcin Nobis<sup>3,4</sup> & Agnieszka Nobis<sup>3</sup>

<sup>1</sup>Department of Biosystematics, Laboratory of Geobotany & Plant Conservation, Opole University, Oleska 22, 45-052 Opole, Poland, anowak@uni.opole.pl; snowak@uni.opole.pl;

<sup>2</sup>Department of Biology and Ecology, University of Ostrava, 71-000 Ostrava, Czech Republic,

<sup>3</sup>Department of Plant Taxonomy, Phytogeography and Herbarium, Institute of Botany, Jagiellonian University, Kopernika 27, 31-501 Kraków, Poland, agnieszka.nobis@uj.edu.pl;

<sup>4</sup>Laboratory of Biodiversity and Ecology, Institute of Biology, Tomsk State University, 36 Lenin Prospect, Tomsk, 634050, Russia, m.nobis@uj.edu.pl

\*Corresponding author

#### **Abstract**

This paper is the continuation of a survey of the scree vegetation in alpine landscapes of western Pamir-Alai Mts in Tajikistan. In total, 105 phytosociological relevés were collected in the colline and montane belt in 2012–2014, applying the Braun-Blanquet method. Phytocoenoses inhabiting mobile or fairly stabilised screes of different sizes of rock particles in the montane and colline zone are herein described. A hierarchical syntaxonomic synopsis of scree communities in the western Pamir Alai Mts is provided. The collected vegetation samples represent the majority of the variations among the phytocoenoses of gravel, pebble, cobble and rock block slides and screes. As a result of field survey and numerical analyses, eight associations – *Cousinietum corymbosae*, *Eremostachyetum tadschikistanicae*, *Cousinietum refractae*, *Caccinietum dubiae*, *Eremuretum sogdiani* (with two subassociations: *E. s. typicum* and *E. s. delphinietosum decolorati*), *Feruletum kuhistanicae*, *Zygophylletum atriplicoidis* and *Corydalidetum kashgaricae* – have been described. Because of their floristic composition, all of these communities have been assigned to a new alliance *Alceion nudiflorae* within the *Sileno brahuicae-Scutellarietalia intermediae* order and *Artemisio santolinifoliae-Berberidetea sibiricae* class. The main factors determining the species composition of the classified associations seem to be scree mobility, rock particle size, elevation above sea level and slope inclination. Saxicolous vegetation in Tajikistan reveal an extraordinary diversity and richness in terms of species composition and beta diversity along the main environmental gradients in this mountainous areas. A further geobotanical survey is needed, especially in eastern Pamir and adjacent areas to fully recognize the chasmophytic plant communities of this rocky land.

**Keywords:** *Artemisio-Berberidetea*, chasmophytes, colluvial cones, phytosociology, Tadjikistan

**Erweiterte deutsche Zusammenfassung am Ende des Artikels**

## 1. Introduction

The landscape of Tajikistan is dominated by mountain ridges and vast areas of screes, taluses, colluvial cones, fans and aprons as well as glaciers and moraines in the upper subnival belt. The Pamir-Alai ranges cross the western part of the country, Tian Shan enter its territory from the north while eastern outskirts are the bareland of the subnival Pamirian Plateau. Almost half of the country's area is made up of bare, rocky and often deserted land dominated by rupicolous habitats. The country is located in Middle Asia, surrounded by ridges of the Karakorum, Hindu Kush, Kunlun and Tian Shan Mts. Because of its considerable biogeographical separateness, diverse climatic conditions and various geology and geomorphology, Tajikistan is very rich in plant species and vegetation types. According to the ten-volume flora of the former Soviet Socialist Republic of Tajikistan, ca 4,550 vascular plant species occur in the country (RASULOVA 1991). This number was supplemented recently by some new records of Tajik flora (e.g. NOBIS 2011a, NOBIS & NOWAK 2011, NOBIS et al. 2014a, b, 2015). Some species new to science have been described from this country in the past few years (e.g. NOBIS 2011b, 2013, NOBIS et al. 2013). Rocky, chasmophytic habitats in Tajikistan, probably because of their high degree of isolation and influences of the Irano-Turanian, Boreal and Indo-Chinese floras, act as a hotspot and refuge for many stenochoric plant species. Approximately 30% of the vascular plant species are considered as endemics of the country. More than half of them inhabit screes and rocks. Scree phytocoenoses also harbour the highest portion of endemics being exclusive to this type of vegetation (NOWAK et al. 2011). Because of this considerable floristic richness, the mountains of Middle Asia have been recognized as being among the world's top 34 biodiversity hotspots (MITTERMEIER et al. 2005), and as one of the 11 most important focal points of scientific studies and conservation efforts (GIAM et al. 2010).

Vegetation of screes has attracted the attention of several geobotanists in recent decades, mainly in Europe and south-western Asia. Studies on mountain colluvial habitats has been conducted in the Carpathians (e.g. KOSIŃSKI 2001, CHYTRÝ 2009, SANDA et al. 2008), the Alps (e.g. ENGLISCH et al. 1993), and mountains of the Mediterranean province (MUCINA et al. 1990, MOTA POVEDA et al. 1991, DIMOPOULOS et al. 1997, DEIL et al. 2008, ORTIZ & RODRIGUEZ-OUBIÑA 1993). For Europe as a whole, a general view of scree vegetation syntaxonomy is presented in the valuable work of VALACHOVIČ et al. (1997). In Africa, only few studies concerning rupicolous vegetation have been published (e.g. DEIL & HAMMOUMI 1997). In south-west Asia, the Taurus (HEIN et al. 1998, PAROLLY 1998, EREN et al. 2004), Alborz (NOROOZI et al. 2014) and Caucasus (ERMOLAEVA 2007, GOLUB et al. 2009, BELONOVSKAYA 2012, BELONOVSKAYA et al. 2014) primarily attracted researchers. However, there is still a lack of comprehensive studies devoted to scree vegetation in Middle (mainly Kyrgyzstan, Tajikistan and Uzbekistan) and Central Asia (NE China, Mongolia; for details, see COWAN 2007), notwithstanding some first insights into this topic presented by ERMAKOV et al. (2006). For a long period, a few contributions to the scree flora of the Hissar Mts (DZHURAEV 1970, 1972a, b) have been the only studies on scree vegetation in Tajikistan. The first phytosociological work based on the Braun-Blanquet method was presented by NOWAK et al. (2015). This research was devoted to the scree vegetation of the alpine and subnival belt.

Despite its enormous uniqueness in terms of floristic diversity Tajikistan still waits for a comprehensive and detailed description of vegetation units and their distribution patterns and habitat characteristic. A geobotanical survey aiming at a precise presentation of Tajik vegetation was initiated in 2006. Segetal and littoral phytocoenoses were the first being

described according to the Braun-Blanquet approach (e.g. NOWAK & NOWAK 2013, NOWAK A. et al. 2013, NOWAK S. et al. 2013a, b). Also for forest and aquatic habitats several syntaxonomic units have been proposed (e.g. NOWAK & NOBIS 2012, 2013). Recently, the petrophytic vegetation of rock faces, crevices and ledges in the alpine and colline zones has been classified (NOBIS et al. 2013, NOWAK A. et al. 2014a, b, c, d).

Despite some floristic similarities between *Asplenietea trichomanis* communities and scree vegetation in Middle Asia, there are some crucial differences in environmental conditions and consequently in the species composition. Thus, in order to establish a complete syntaxonomic system for the rupicolous vegetation of Tajikistan, the scree of colluvial deposits must be studied in detail. This paper presents the first attempt to classify the scree vegetation inhabiting the colline and montane belts of Pamir-Alai. We aimed at addressing the following questions in our study: (1) What is the diversity of the scree vegetation of colline and montane zones in the Pamir Alai Mts? (2) What are the habitat conditions of the described plant communities? (3) What is their species composition and structure? (4) Which species have important diagnostic value for the described syntaxa?


Considering our former studies devoted to rock vegetation (e.g. NOWAK A. et al. 2014a, b) the final aim of the project is to establish a syntaxonomic classification of all chasmophytic habitats of Middle Asia (at least Pamir-Alai and Tian Shan Mts).

## 2. Methods

### 2.1. Study area

Tajikistan is located in Middle Asia between 36°40'–41°05' E and 67°31'–75°14' N and covers ca 143,500 km<sup>2</sup> (Fig. 1). It is a mountainous country, with more than 50% of the area located above 3,000 m and more than 93% above 1,000 m a.s.l. As part of the Eurasian highland belt, Tajikistan encompasses a vast area covered by stony wasteland with diversified colluvial and diluvial deposits in terms of exposition, inclination, vertical zonation and particle size (blocks, cobbles, pebbles, coarse and fine gravels). The southern part of Tajikistan is influenced by a subtropical climate, while the northern part is situated in the temperate climate zone (VLADIMIROVA 1968). Generally, the area is characterised by high solar radiation (2,090–3,160 sunshine hours), a low percentage of cloud cover, a high amplitude of annual temperatures, low humidity and low precipitation. In subtropical regions of Tajikistan, average temperatures in June are around 30 °C. In the temperate zone, in Tajikistan comprising mainly the high mountains, the climate is much harsher, with average temperatures in July between 9.7 °C and 13.5 °C. Annual precipitation ranges from ca. 70 mm in the eastern Pamir to ca 600 mm in the Hissar Range (in some locations up to ca. 2,000 mm). In the western part of the country, the lower limit of perpetual snow is at an altitude of 3,500–3,600 m a.s.l.; in its eastern regions, at 5,800 m (LATIPOVA 1968, NARZIKULOV & STANJUKOVICH 1968, SAFAROV 2003).

The investigated area falls generally into the colline and montane zones of several mountain ranges with typical rugged topography: the Zeravshan, Hissar, Hazratishoh, Darvaz, Vanch, Turkestan, Peter I and Karateginian Mts (Fig. 2). All of them belong to the Pamir-Alai mountain system, which stretches westwards from the Pamir plateau. Only few sample plots were located in the hilly landscape of southwestern Tajikistan or reach the alpine belt in Zeravshan Mts. Alongside all mountain ridges and walls, a considerable number of colluvial cones have developed, generally in the bottoms of the V-shaped valleys of streams and brooks, due to frost erosion and gravitational forces. These cones consist of different types of heterogeneous and generally unsorted parent material that have moved down the slope. Rock particles are sharp and angular and of various sizes (ranging from blocks, cobbles, pebbles, and coarse- and fine-grain gravel to sandy silt and clay). The studied scree differs also in terms of exposition, inclination, bedrock type and altitude. The studied vegetation plots were located between 359 m and 2,420 m a.s.l. (mean ca 1,600) on rock substrates with pH ranging from 6.5 to 8.8.




**Fig. 1.** The location of Tajikistan in Middle Asia.

**Abb. 1.** Die Lage Tadschikistans in Mittelasien.

The geological structure of the study area is very complex, with outcrops of rocks formed from the Precambrian to the present age. Only few geological studies have been published for Tajikistan (e.g. NEDZVEDSKIY 1968). The middle and higher parts of the Hissar Mts are largely composed of extrusive rocks, mainly granite, granitoid and syenite. Some igneous outcrops also occur in the Darwaz and Kuraminian Mts and in the western Pamir ranges. In the Zeravshan and Turkestan Mts, Cambrian and Silurian sediments predominate. The rocks here are generally limestone (micritic, bitumic, marly and dolomitic coral limestone), dolomite, dolomitic shale clay shale and conglomerates. Several kinds of metamorphic rocks are also present within the study area. The most common are migmatic gneiss, marble, phyllitic schists, argillaceous slates and metamorphic mudstones.

## 2.2. Data sampling and analyses

The main research season was conducted in 2014, however, several samples were also gathered in 2010, 2011 and 2012. Approximately 250 relevés were collected in screes in the Pamir-Alai Mts with 105 in colline and montane belts. The montane zone extends in western Pamir-Alai system from approx. 900 m to 2,000–2,500 m a.s.l. depending on mountain range and slope exposition. The upper limit of this belt is related to the tree line and is generally marked by juniper forests (*Juniperus turkestanica*, *J. zeravshanica*). The colline zone starts from ca 400 m a.s.l. and is mainly characterized by azonal riverside forests of *Populus pruinosa*, grasslands, thermophilous swards and shrubs. The plot size was generally 30 m<sup>2</sup> except for communities with *Corydalis kashgarica* where a plot size of 5 m<sup>2</sup> was applied due to different community physiognomy. In each relevé, all vascular plant species were recorded using the seven-degree cover-abundance scale of Braun-Blanquet (WESTHOFF & VAN DER MAAREL 1973). Cryptogams were not recorded because of their negligible abundance and importance in the scree communities in the study area. The aim of the survey was to cover a broad range of habitats in relation to altitudinal range, size of rock debris, exposition and inclination. Following the field research, we selected 105 relevés to present the scree vegetation of the colline and montane zone in western Tajikistan. For each plot, geographical coordinates were measured with the help of a GPSMAP 60CSx



**Fig. 2.** Topographic characterisation of the study area with main mountain ranges, rivers and cities.

**Abb. 2.** Topographischer Überblick des Untersuchungsgebiets mit den wichtigsten Gebirgszügen, Flüssen und Städten.

device with an accuracy of  $\pm 5$  m, using the WGS84 reference frame. In the tables, latitude and longitude are given in degrees, minutes and seconds. Rock type was determined through an analysis of lithology, pore geometry, mineralogical components, texture, permeability, hardness and pH, performed by a professional geologist (see acknowledgments). Hydrogen ion concentrations were measured in an aqueous rock solution using an Elmetron CP-105 pH meter.

Data were stored in the JUICE program (TICHÝ 2002). A modified TWINSpan classification (ROLEČEK et al. 2009) provided an initial idea of data structure and resolution. We applied pseudospecies cut levels of 0%, 2%, 5% and 10%. A Detrended Correspondence Analysis (DCA) was performed using the floristic data set (no downweighting of rare species) to check the floristic-sociological classification and to highlight the relationship between groups. CANOCO for Windows 4.5 was used for this ordination (TER BRAAK & ŠMILAUER 2002). The species composition data showed a clear unimodal response, with a total gradient length of ca 9 for the complete data set.

Vegetation classification followed the sorted table approach of BRAUN-BLANQUET (1964). A synoptic table with the percentage frequencies of diagnostic species is given in Table 1. For newly-described associations, the International Code of Phytosociological Nomenclature was adhered to (WEBER et al. 2000). The class concept follows PIGNATTI et al. (1995), with general habitat features and a common set of geographically distinct character species as key traits defining the class. As the most significant attributes of the habitat, we considered spatial structure and environmental characteristics, mainly the presence of moving rock or gravel deposits creeping down the slope. When analysing the possible class assignment of the associations, we considered mainly the *Thlaspietea rotundifolii* Br.-Bl. 1948, as a syntaxon comprising the scree vegetation of Europe; the *Heldreichietea* Quezel ex Parol. 1995, which include scree vegetation of the east Mediterranean region; the *Lamio tomentosio-Chaerophylletea humilis* Belonovskaya, Mucina et Theurillat 2014 reported from screes in Caucasus

(BELONOVSKAYA et al. 2014) and the *Artemisia santolinifoliae-Berberidetea sibiricae*, which have been proposed for vegetation types of scree and rocky slopes of Central Asia (ERMAKOV et al. 2006). All of these syntaxa are climax, stable vegetation types inhabiting extreme environments, with relatively few species per plot.


Species nomenclature followed mainly CHEREPANOV (1995) and the Plant List (<http://www.theplantlist.org>; accessed 15 May 2015). Plant material collected during field studies was deposited in the Herbarium of Middle Asia Mountains, hosted in OPUN (Opole University, Poland) and KRA (Jagiellonian University, Poland).

### 3. Results

The number of taxa recorded in the relevés totals 198, with 78 exceeding 5% constancy and 42 exceeding 10%. To the group of vascular plant species with highest frequency belong: *Bromus tectorum* (54 occurrences), *Centaurea squarrosa* (54), *Alcea nudiflora* (47), *Trichodesma incanum* (41), *Artemisia persica* (32), *Pulicaria salvifolia* (29), *Origanum tyttanthum* (26), *Piptatherum songaricum* (26), *Callipeltis cucullaris* (25), *Sanguisorba alpina* (24), *Salvia sclarea* (20), *Galium spurium* (19), *Glaucium elegans* (19), *Hypericum scabrum* (19), *Scariola orientalis* (19) and *Scrophularia scabrosifolia* (19). Several species contributing to the surveyed plant communities are endemics of the Pamir-Alai, restricted in most cases to the territory of Tajikistan, e.g. *Eremostachys mogianica*, *E. tadschikistanica*, *Allium crystallinum*, *Astragalus varzobicus*, *Cousinia butkovii*, *C. corymbosa*, *C. outichaschensis*, *C. pseudohisakensis*, *C. refracta*, *Delphinium decoloratum*, *Eremurus comosus*, *E. olgae*, *E. pubescens*, *Ferula equisetacea*, *F. koso-polianskyi*, *F. sumbul* and others. The extreme uniqueness of the scree flora of the study area (NOWAK et al. 2011) results in almost no floristic affinities with European or south-western Asian mountainous regions. Among the diagnostic species no plants occurring both in Pamir-Alai and those regions were found. Even at the genus level, only few are common to the scree vegetation of Middle Asia and SE Asia or Europe (e.g. *Papaver* sp., *Poa* sp., *Allium* sp., *Veronica* sp., *Cerastium* sp.; VALACHOVIĆ et al. 1997, BELONOVSKAYA et al. 2014).

Not all of the species noted in relevés are typical for scree vegetation. In the studied plots some ruderal species or plants typical for nitrophilous and often anthropogenic habitats were noted, sometimes with considerable abundance and frequency: *Bromus tectorum*, *Poa bulbosa*, *Descurainia sophia*, *Capparis spinosa*, *Koelpinia linearis*. Species typical for agrocenoses have also been noted (*Avena trichophylla*, *Setaria pumila*, *S. viridis*, *Strigosella africana*, *Vaccaria hispanica* or *Lactuca serriola*) as well as plants typical of xerothermophilous swards (e.g. *Hordeum bulbosum*, *Pimpinella puberula*, *Handelia trichophylla*, *Achillea biebersteinii*, *Elytrigia trichophora* or *Aphanopleura capillipes*). A few tall herb species enter the scree communities of Tajikistan (e.g. *Tragopogon krashennikovii*, *Ligularia thompsoni*, *Prangos pabularia* or *Rumex paulsenianum*), forest plants (*Poa pratensis*) or species typical for steppe vegetation (*Erodium oxyrhynchum*). Astonishingly, no species with the optimum in rocky habitats were recorded, despite the insignificant occurrence of *Artemisia rutifolia*, *Carex koshevníkovii* and *Poa relaxa*.

As a result of the numerical classification, 11 groups were distinguished (Fig. 3). The two classes presented in the syntaxonomic scheme correspond to the first division in the TWINSPLAN classification. The cluster 1 comprised the *Corydalisetum kashgaricae* of the *Parietarietea judaicae* class. The cluster 0 accommodated all other communities. It was divided into cluster 0–1, which included the driest habitats (*Zygophylletum atriplicoides* and community with *Echinops nanus*), and cluster 0–0, representing the moderately humid, typi-




**Fig. 3.** DCA ordination of all samples of *Alceion nudiflorae* communities ( $n = 105$ ).

**Abb. 3.** DCA-Ordination aller Aufnahmen des Verbands *Alceion nudiflorae* ( $n = 105$ ).

cal screens of the montane zone. The third level of division was mainly related to substrate acidity. Within cluster 0–0–1, communities preferring strongly alkaline habitats were grouped (*Eremuretum sogdianae*, *Caccinietum dubiae* and community of *Glaucium elegans*). Cluster 0–0–0 was mainly composed of relevés from calcifuge substrates (*Cousinietum corymbosae*, *Eremostachyetum tadschikistanicae* and *Cousinietum refractae*). This distinction according to alkalinity, however, was not so clear because the last cluster included also the calcicole *Feruletum kuhistanicae* and community with *Zygophylletum gontscharovii*. Both associations were separated from the calcifuge ones at the fourth level of division.

According to habitat differences and floristic distinctiveness, eight groups have been identified as associations and defined by original diagnosis based on species composition. Within the calcifuge group, which occupies the central part of the DCA graph (Fig. 3), *Cousinietum corymbosae*, *Eremostachyetum tadschikistanicae* and *Cousinietum refractae* have been distinguished. The group of calcicole communities consists of *Caccinietum dubiae*, *Eremuretum sogdiani* (with subassociation of *delphinietosum decoloratae*), *Feruletum kuhistanicae*, *Zygophylletum atriplicoidis* and *Corydalidetum kaschgaricae*. On the ordination graph, also plots dominated by *Echinops nanus*, *Glaucium elegans* and *Zygophyllum gontscharovii* are shown (Fig. 3). Because of scarcity of data and doubts about their development stage or habitat affinity, these groups have only been assigned as communities with-



**Fig. 4.** Box-whisker plots of the species richness, cover of the herb layer, altitudinal distribution, inclination, pH and size of the rock particles of screes of the researched communities: *Eremuretum sogdiani* (Ere so), *Caccinietum dubiae* (Cac du), *Feruletum kuhistanicae* (Fer kuh), *Zygophylletum atriplicoidis* (Zyg at), *Cousinietum corymbosae* (Cou co), *Eremostachyetum tadschikistanicae* (Ere ta), *Cousinietum refractae* (Cou re), communities of *Glaucium elegans* (Gla el), *Echinops nanus* (Ech na), *Zygophyllum gescharovii* (Zyg go) and the *Corydalidetum kashgaricae* (Cor ka).

**Abb. 4.** Box-Whisker-Plots mit Artendiversität, Deckung der Krautschicht, Seehöhe, Hangneigung, pH-Wert und Größe der Gesteinsbruchstücke in den untersuchten Gesellschaften: *Eremuretum sogdiani* (Ere so), *Caccinietum dubiae* (Cac du), *Feruletum kuhistanicae* (Fer kuh), *Zygophylletum atriplicoidis* (Zyg at), *Cousinietum corymbosae* (Cou co), *Eremostachyetum tadschikistanicae* (Ere ta), *Cousinietum refractae* (Cou re), *Glaucium elegans*-Gesellschaft (Gla el), *Echinops nanus*-Gesellschaft (Ech na), *Zygophyllum gescharovii*-Gesellschaft (Zyg go) und *Corydalidetum kashgaricae* (Cor ka).



out rank. The same is true for plots with *Chondrilla gibbistrois*, which forms rather mono-specific stands along road verges, and *Prangos pabularia*, which has its ecological optimum in forb vegetation. Further research is needed to find the final syntaxonomic position of these communities.

The habitat characteristics and species composition suggests to classify most of the phytocoenoses into the class *Artemisio santolinifoliae-Berberidetea sibiricae*, which includes shrubby, species poor vegetation on mobile scree in Central Asia. This syntaxonomic assignment is justified by comparison of the life form proportions within particular plots. As in communities of the *Artemisio-Berberidetea* (ERMAKOV et al. 2006), we observed a considerable share of dwarf shrubs and shrubs. Small shrubs like *Artemisia* spp., *Krashennikovia ceratoides* or *Gontscharovia popovii* had a absolute cover up to 16% (average approx. 3%) and higher shrubs (e.g. *Rosa* spp., *Berberis* spp., *Ephedra* spp., *Cerasus verrucosus* or *Atraphaxis pyrifolia*) reached an average absolute cover of approx. 4% (maximum 40%) in our plots. However, in all vegetation types the perennials clearly dominated (approx. 80% on average) with some contribution of annuals and hemicryptophytes. Considerably different in terms of floristic composition was the *Corydalidetum kashgaricae*. The occurrence of nitrophilous species and the generally low cover of the herb layer are important features of that type vegetation. These plots were classified into the *Parietarietea judaicae* Oberd. 1977.

#### 4. Description of the syntaxa

We propose the following classification of the colline and montane scree vegetation of the Pamir-Alai Mts in Tajikistan:

Class: *Artemisio santolinifoliae-Berberidetea sibiricae* Ermakov et al. 2006

Order: *Sileno brahuicae-Scutellarietalia intermediae* nom. prov.

Alliance: *Alceion nudiflorae* all. nova

Calcifuge group:

1. *Cousinietum corymbosae* ass. nova
2. *Eremostachyetum tadschikistanicae* ass. nova
3. *Cousinietum refractae* ass. nova

Calcicole group:

4. *Caccinietum dubiae* ass. nova
5. *Eremuretum sogdiani* ass. nova
  - 5.1. *E. s. typicum*
  - 5.2. *E. s. delphinietosum decolorati* subass. nova
6. *Feruletum kuhistanicae* ass. nova
7. *Zygophylletum atriplicoidis* ass. nova
8. Community with *Echinops nanus*
9. Community with *Glaucium elegans*
10. Community with *Prangos pabularia*
11. Community with *Zygophyllum gontscharovii*
12. Community with *Chondrilla gibbistrois*

Class: *Parietarietea judaicae* Oberd. 1977

Order: *Tortulo-Cymbalarietalia* Segal 1969

Alliance: *Parietarion judaicae* Segal 1969

13. *Corydalidetum kashgaricae* ass. nova

### **Alliance: *Alceion nudiflorae* all. nova**

Nomenclatural type: *Eremuretum sogdiani* holotypus hoc loco

Diagnostic species: *Alcea nudiflora*, *Trichodesma incanum*, *Piptatherum songaricum*, *Pulicaria salvifolia*

Distribution and ecology: The *Alceion nudiflorae* alliance was recorded in western Pamir-Alai Mts, mainly in Zeravshan, Hissar, Hazratishoch, Peter I, Turkestan, Darwaz and Vachsh ranges. It occurs at (350–)1,200–1,900(–2,400) m a.s.l. The phytocoenoses of *Alceion nudiflorae* are moderately rich in species with somewhat higher species number per plot compared to the alpine and subnival scree phytocoenoses of the *Ferulo foetidissimae-Vicion kokanicae* (NOWAK et al. 2015). The communities are developed on various rock substrata (generally limestone, marble, dolomite, schists and granite) of different size (cobbles, pebbles, coarse and fine grained gravel). A considerable portion of species composing the alliance have the Irano-Turanian distributional type. All diagnostic taxa are distributed throughout Middle Asia (generally Pamir-Alai and Tian Shan Mts), Iran, northern Afghanistan and Pakistan, western Himalayas and China (KOCHKAREVA 1986, OVCHINNIKOV 1957, KINZIKAEVA 1988). Apart from the taxa diagnostic for the alliance, scree species with higher frequencies are *Artemisia persica*, *Callipeltis cucullaris*, *Sanguisorba alpina*, *Salvia sclarea*, *Hypericum scabrum*, *Lactuca orientalis* and *Scrophularia scabiosifolia* (Tables 1–3).

#### **1. *Cousinietum corymbosae* ass. nova**

Typus relevé: Supplement S1, rel. 2, holotypus hoc loco

Diagnostic species: *Cousinia corymbosa*, *Ferula equisetacea*


The phytocoenoses of *Cousinietum corymbosae* have been found in several locations in Hissar Mts, mainly on western and south-western slopes in Varzob, Takob and Romit River Valleys (Fig. 6, 7). *Cousinia corymbosa* is a narrow endemic of Hissar Mts, scattered on few locations near Maychura, Varzob, Khokhu and Romit villages on gravel deposited at hillslope base, at elevations 2,000–2,400 m a.s.l. (RASULOVA 1991). The altitudinal distribution of the association is not fully compliant with this amplitude. The samples have been taken in colline and montane zone at the elevation from 1,230 to 1,850 m a.s.l. (mean approx. 1,600; Fig. 4, Supplement S1). The association prefers granite, relatively loose and fine-grained screes (pH 6.5–7.4). The *Cousinietum corymbosae* was found mainly on western and south-western expositions with inclinations of mean value approx. 35° (Fig. 5). The association is characterised by a moderate vegetation cover. The total cover of the herb layer was close to 60% on average (Supplement S1, Fig. 4). The phytocoenosis is characterised by a moderate number of species as scree vegetation is concerned, having in one relevé from 10 to 23 taxa (mean value approx. 13). Among the vascular plants the highest values of constancy and abundance within sampled plots have: *Alcea nudiflora*, *Bromus tectorum*, *Callipeltis cucullaris*, *Hypericum scabrum*, *Impatiens parviflora*, *Origanum tyttanthum*, *Scrophularia scabiosifolia*, *Sanguisorba alpina* and *Piptatherum songaricum*.

#### **2. *Eremostachyetum tadschikistanicae* ass. nova**

Typus relevé: Supplement S1, rel. 14, holotypus hoc loco


Diagnostic species: *Eremostachys tadschikistanica*

*Eremostachys tadschikistanica* is a narrowly distributed rupicolous species, considered as endemic of Hissar Mts (KOCHKAREVA 1986). During our studies the association of *Eremostachys tadschikistanica* was found in several locations along the Takob and Honaka



**Fig. 5.** The exposition preferences of the researched plant communities.  
**Abb. 5.** Die Expositionsverteilung in den untersuchten Gesellschaften.

Rivers on southern faces of Hissar Range (Fig. 6). The phytocoenosis have been found mainly on granite and schist screes with some admixture of dolomites creeping downslope from upper rock ridges (pH 6.0–7.2). The association inhabits screes with fine- or coarse gravels with insignificant soil amount. The phytocoenosis develops on relatively high elevations within the range of montane belt (mean ca. 1,800 m a.s.l) with some plots entering also alpine zone (Fig. 4). The community develops on fairly steep screes with mean inclination of approx. 45°. *Eremostachysetum tadschikistanicae* is strictly related to southern expositions with very high solar radiation. The mean cover of the herb layer is almost the highest among the montane scree communities with about 65%, ranging from 60 to 80% (Fig. 4, Supplement S1). The most abundant and constant vascular plant species are: *Alcea nudiflora*, *Bromus danthoniae*, *B. tectorum*, *Callipeltis cucullaris*, *Centaurea squarrosa*, *Eremurus comosus*, *Hypericum scabrum*, *Origanum tyttanthum*, *Sanguisorba alpina*, *Trichodesma incanum* and *Veronica intercedens*.



Explanations: I: ● *Cousinietum corymbosae*, ■ *Caccinietum dubiae*, △ *Eremuretum sogdiani*, II: ◆ *Eremostachyetum tadschikistanicae*, ○ *Cousinietum refractae*, ◊ *Feruletum kuhistanicae*, III: □ *Zygophyllum atriplicoides*, ● *Comm. with Zygyphyllum gontscharovii* IV: ◊ *Comm. with Glaucium elegans*, ▲ *Comm. with Chondrilla gibbrosistris*, ✚ *Comm. with Echinops nanus*, ✚ *Comm. with Prangos pabularia*

**Fig. 6.** Distribution of the associations of montane and colline screes in Tajikistan.


**Abb. 6.** Verbreitung der Assoziationen der montanen und kollinen Schuttvegetation in Tadschikistan.

### 3. *Cousinietum refractae* ass. nova

Typus relevé: Supplement S1, rel. 21, holotypus hoc loco

Diagnostic species: *Cousinia refracta*

*Cousinia refracta* is a stenochoric species with a range restricted to screes and slopes in south-western Tajikistan, mainly in Hissar, Aktau and Karatau Mts (RASULOVA 1991). The association was found in the montane zone on the warmest slopes in close contact with xerothermophilous swards and shrubs. The association develops in Honaka and Khondara River valleys and near Gusgarf (Fig. 6). It was found exclusively on granite screes of moderately compact structure (pH 6.4–7.5). In comparison to other plant communities a considerable amount of clayey soils were noted in research plots. The rock particles were relatively fine and compact here. The phytocoenosis develops on relatively low elevations in the colline zone with warm microclimate. The observed altitudinal range was between 1,350 and 1,650 m a.s.l. (mean approx. 1,600). The community develops on moderately steep screes with the mean inclination value for the samples of approx. 35°, ranging from 30° to 50° (Fig. 4). *Cousinietum refractae* inhabits generally south-western expositions (Fig. 5). The mean cover of the herb layer was among the highest and was close to 70%, ranging from 45 to 80% (Fig. 4). The plots of vegetation were moderately rich in vascular species. As a co-dominant species within the sampled plots *Alcea nudiflora*, *Centaurea squarrosa* and *Salvia*



**Fig. 7.** (a) *Caccinietum dubiae* on a pebble steep scree near Zerabad in Zeravshan Mts; (b) *Cousinietum refractae* on a granite fine scree in Varzob valley in Hissar Mts; (c) *Zygophylletum atriplicoidis* on southern slopes of Turkestan Mts in Zeravshan River Valley near Zasan; (d) *Feruletum kuhistanicae* on apron cobbles in Seven Lake valley near Nofin; (e) plot with dominance of *Glaucium elegans* on a fine gravel scree near Sarytog; (f) community with *Zygophyllum gontscharovii* in south-western Tajikistan near Garavuti.

**Abb. 7.** (a) *Caccinietum dubiae* auf einer steilen Geröllhalde nahe Zerabad im Serafschan-Gebirge; (b) *Cousinietum refractae* auf einer feinen Granithalde im Varzob-Tal im Hissargebirge; (c) *Zygophylletum atriplicoidis* auf südexponierten Hängen der Turkestankette im Tal des Serafschan bei Zasan; (d) *Feruletum kuhistanicae* auf Vorfeld-Blöcken im Sieben-Seen-Tal bei Nofin; (e) Fläche mit Dominanz von *Glaucium elegans* auf einer Feinkies-Halde nahe Sarytog; (f) Gesellschaft mit *Zygophyllum gontscharovii* in Südwest-Tajikistan bei Garavuti.

*sclarea* were found. Other fairly frequent species contributing to this association were *Aegilops triuncialis*, *Astragalus varzobicus*, *Avena trichophylla*, *Eremurus olgae*, *Hordeum bulbosum*, *Pimpinella puberula*, *Sanguisorba alpina* and *Trichodesma incanum*.

#### 4. *Caccinietum dubiae* ass. nova

Typus relevé: Supplement S1, rel. 31, holotypus hoc loco

Diagnostic species: *Caccinia dubia*

*Caccinia dubia* is a species of broader distribution in Middle Asia, known from screes of Pamir-Alai and Tian Shan Mts (Fig. 1, 7). In Tajikistan the species is known from northernmost ranges (Kuraminian, Turkestan and Zeravshan) from montane and lower alpine elevations (CHUKAVINA 1984; Fig. 6). The association was found mainly on schists and shales of alkaline reaction (pH 8.0–8.6). However, field observations and literature confirms also occurrence of *Caccinia dubia* on neutral or acidic substrates (CHUKAVINA 1984). The association prefers screes almost deprived of any soil ingredient with relatively large rock pebble and cobble colluvial slides (Fig. 4). The community inhabits the driest areas at moderate elevations in montane zones, within the altitudinal range between 1,400 and 2,050 m a.s.l. (mean approx. 1,700). Plots of *Caccinietum dubiae* develops on fairly steep screes, with the mean inclination value approx. 45°, ranging from 35° to 55°. The phytocoenosis inhabits generally southern expositions (Fig. 4, 5). The cover of the herb layer is relatively low, ranging from 30 to 55% (mean approx. 40%; Fig. 5). The most abundant and constant species are: *Alcea nudiflora*, *Bromus tectorum*, *Centaurea squarrosa*, *Cousinia ferganensis*, *Echinops chloroleucus*, *Nepeta olgae*, *Pulicaria salvifolia*, *Lactuca orientalis* and *Trichodesma incanum*.

#### 5. *Eremuretum sogdiani* ass. nova

Typus relevé: Supplement S1, rel. 43, holotypus hoc loco

Diagnostic species: *Eremurus sogdianus*

*Eremurus sogdianus* is commonly distributed along the foothills of south-western Tian Shan and northern Pamir-Alai. Only in Zeravshan Mts the species enters the northern valleys more deeply (OVCHINNIKOV 1963, KOVALEVSKAYA 1971). The distribution of the association is restricted to lower sections of northern valleys of Turkestan and Zeravshan ranges (Fig. 6). The research conducted in this area confirms that the species builds its own association developing on alkaline rock debris (pH 7.7–8.8), mainly on dolomite and limestone screes. The community inhabits colluvial deposits with coarse gravels with mean particle size of approx. 4 cm (Fig. 4). The association prefers elevations in montane zone (Fig. 4), within the altitudinal range between 1,200 and 1,800 m a.s.l. (mean approx. 1,600). The community develops on gently sloping screes with mean inclination close to 30°; Fig. 4). Unlike other plant communities from the *Alceion nudiflorae*, *Eremuretum sogdiani* prefers generally north-western and western, somewhat shadowed expositions (Fig. 5). Within the sampled plots, between 10 and 24 taxa were noted (mean approx. 16), so as the majority of the rupicolous vegetation in Tajikistan, the association should be classified as moderately rich in species. Also the cover of the herb layer is moderate and reaches up to 75% with the mean value of approx. 60% (Fig. 5). Among the species occurring within the association with higher constancies are *Alcea nudiflora*, *Artemisia persica*, *Bromus tectorum*, *Cousinia radians*, *Crambe kotschyana*, *Koelpinia linearis*, *Perovskia virgata*, *Lactuca orientalis*, *Scutellaria intermedia* and *Trichodesma incanum*.

### 5.1. *Eremuretum sogdiani typicum* subass. nova

Type relevé: Supplement S1, rel. 43, holotypus hoc loco

Diagnostic species: *Eremurus sogdianus*

Constant species: the same as for the association

The subassociation *typicum* is characterised by higher share of xerothermophilous sward species and slightly more arid microhabitat. It seems that the *E. s. typicum* prefers sites with long and intensive solar radiation on gentle slopes of wide valleys. The range of the typical subassociation encompass whole Middle Asia. In comparison to *Eremuretum sogdiani delphinietosum decolorati* plots of subassociation *typicum* are more frequently entered by *Alyssum desertorum*, *Cousinia ferganensis*, *Taeniatherum crinitum*. The habitat characteristic, diagnostic species and floristic composition of this subassociations are the same as for the association.

### 5.2. *Eremuretum sogdiani delphinietosum decolorati* subass. nova

Type relevé: Supplement S1, rel. 54, holotypus hoc loco

Diagnostic species: *Delphinium decoloratum*

This subassociation was found sporadically in the lower section of the Seven Lake valley in the western part of Zeravshan Mts (Fig. 6). The association is distributed in the montane zone at moderate altitudes from 1,650 to 1,770 m (Fig. 4). It prefers coarse-grained, moderately compact gravel screes. The community was observed on limestone debris with pH 8.0–8.4. The subassociation occurs generally on eastern and northern expositions with sloping rate of approx. 30°–45° (Fig. 4, 5). *Eremuretum sogdiani delphinietosum decolorati* inhabits shadowed sites in narrow gorges of Zeravshan range. The total cover of the herb layer was moderate, between 50 and 65%. The phytocoenosis is characterised by fairly high species richness with the mean value of ca. 23. Among the vascular plants contributing to the herb layer, apart from the diagnostic species, the most constant and abundant are e.g.: *Bromus tectorum*, *Impatiens parviflora* and *Reseda lutea*.

## 6. *Feruletum kuhistanicae* ass. nova

Type relevé: Supplement S2, rel. 5, holotypus hoc loco

Diagnostic species: *Ferula kuhistanica*

The main diagnostic species of the association commonly occurs in Tajikistan except the northern outskirts of the country (CHUKAVINA 1984). *Ferula kuhistanica* grows also in Tian Shan and northern Afghanistan. The plots of *Feruletum kuhistanicae* were found in several locations in the valleys of the Varzob, Takob, Sarday-Myena and Obihingou River valleys, and in Seven Lake valley (Fig. 6, 7). The community was found in the montane belt at moderate altitudes from 1,250 to 1,800 m a.s.l. (mean approx. 1,500 m). The association inhabits dolomite, limestone and schist screes with pH 8.3–8.8. The deposits are generally fairly mobile, however also plots on compact debris with soil layer were found. The community develops mainly on south-western and southern aspects of gently sloping colluvial cones and aprons built by large stones and cobbles (Fig. 4). The approximate mean inclination was ca. 35°, ranging from 15° to 40°. The plots of *Feruletum kuhistanicae* are relatively rich in species (mean 19 taxa per plot). The association is characterised by a moderate cover of the herb layer, ranging between 50 and 80%, with mean value of approximately 60% (Supple-

**Table 1.** Synoptic table of the scree communities of the montane and colline zones in western Pamir-Alai Mts, Tajikistan. Values are percentage constancies. Only diagnostic species are shown. 1: *Cousinietum corymbosae*; 2: *Eremostachyetum tadschikistanicae*; 3: *Cousinietum refractae*; 4: *Caccinietum dubiae*; 5: *Eremuretum sogdiani*; 6: *Eremuretum sogdiani delphinietosum decolorati*; 7: *Feruletum kuhistanicae*; 8: *Zygophylletum atriplicoidis*; 9: *Corydalidetum kashgaricae*.

**Tabelle 1.** Synoptische Tabelle der montanen und kollinen Schuttvegetation in Tadschikistan. Die Zahlen geben die prozentuelle Stetigkeit an. Nur diagnostische Arten sind dargestellt. Syntaxa s. o.

| Syntaxonomical unit                                     | 1   | 2   | 3   | 4   | 5   | 6   | 7   | 8   | 9  |
|---|-----|-----|-----|-----|-----|-----|-----|-----|----|
| Number of relevés                                       | 10  | 10  | 10  | 11  | 10  | 4   | 11  | 10  | 9  |
| Av. species number per relevé                           | 15  | 20  | 16  | 15  | 15  | 23  | 19  | 8   | 4  |
| TWINSPAN level 1  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1  |
| TWINSPAN level 2  | 0   | 0   | 0   | 0   | 0   | 0   | 0   | 1   |    |
| TWINSPAN level 3  | 0   | 0   | 0   | 1   | 1   | 1   | 0   | 0   |    |
| TWINSPAN level 4  | 1   | 1   | 1   |     |     |     | 0   |     |    |
| <b>Cousinietum corymbosae</b>                           |     |     |     |     |     |     |     |     |    |
| <i>Cousinia corymbosa</i>                               | 100 | .   | .   | .   | .   | .   | .   | .   | .  |
| <i>Ferula equisetacea</i>                               | 30  | .   | .   | .   | .   | .   | 9   | .   | .  |
| <b>Eremostachyetum tadschikistanicae</b>                |     |     |     |     |     |     |     |     |    |
| <i>Eremostachys tadschikistanica</i>                    | .   | 100 | .   | .   | .   | .   | 18  | .   | .  |
| <b>Cousinietum refractae</b>                            |     |     |     |     |     |     |     |     |    |
| <i>Cousinia refracta</i>                                | .   | .   | 100 | .   | .   | .   | .   | .   | .  |
| <b>Caccinietum dubiae</b>                               |     |     |     |     |     |     |     |     |    |
| <i>Caccinia dubia</i>                                   | .   | .   | .   | 100 | .   | .   | .   | .   | .  |
| <b>Eremuretum sogdiani</b>                              |     |     |     |     |     |     |     |     |    |
| <i>Eremurus sogdianus</i>                               | .   | .   | .   | 45  | 100 | 75  | .   | .   | .  |
| <b>Eremuretum sogdiani delphinietosum decolorati</b>    |     |     |     |     |     |     |     |     |    |
| <i>Delphinium decoloratum</i>                           | .   | .   | .   | .   | .   | 100 | .   | .   | .  |
| <b>Feruletum kuhistanicae</b>                           |     |     |     |     |     |     |     |     |    |
| <i>Ferula kuhistanica</i>                               | .   | .   | .   | 9   | .   | .   | 100 | .   | .  |
| <b>Zygophylletum atriplicoidis</b>                      |     |     |     |     |     |     |     |     |    |
| <i>Zygophyllum atriplicoides</i>                        | .   | .   | .   | .   | .   | .   | .   | 100 | .  |
| <b>Alceion nudiflorae</b>                               |     |     |     |     |     |     |     |     |    |
| <i>Alcea nudiflora</i>                                  | 60  | 60  | 80  | 82  | 90  | 75  | 45  | .   | .  |
| <i>Trichodesma incanum</i>                              | .   | 100 | 60  | 73  | 80  | 50  | 64  | .   | .  |
| <i>Pulicaria salvifolia</i>                             | 30  | 20  | .   | 82  | 30  | 50  | .   | 50  | 33 |
| <i>Piptatherum songaricum</i>                           | 80  | 40  | .   | 27  | 30  | 75  | 45  | .   | .  |
| <i>Chondrilla gibbistrostris</i>                        | .   | .   | .   | 9   | .   | .   | 9   | .   | 44 |
| <b>Sileno brahuicae-Scutellarietalia intermediae</b>    |     |     |     |     |     |     |     |     |    |
| <i>Scutellaria intermedia</i>                           | .   | .   | .   | .   | 70  | 75  | 36  | .   | .  |
| <i>Silene brahuica</i>                                  | .   | .   | .   | .   | .   | .   | 36  | .   | .  |
| <b>Artemisio santolinifoliae-Berberidetea sibiricae</b> |     |     |     |     |     |     |     |     |    |
| <i>Centaurea squarrosa</i>                              | 40  | 80  | 80  | 91  | 70  | 75  | 27  | 40  | 56 |
| <i>Artemisia persica</i>                                | 10  | 50  | .   | 18  | 80  | 75  | 45  | .   | .  |
| <i>Callipeltis cucullaris</i>                           | 50  | 80  | .   | 9   | .   | .   | 73  | 10  | 11 |
| <i>Sanguisorba alpina</i>                               | 70  | 60  | 90  | .   | .   | .   | 18  | .   | .  |
| <i>Hypericum scabrum</i>                                | 50  | 70  | 70  | .   | .   | .   | .   | .   | .  |
| <i>Salvia sclarea</i>                                   | 10  | 20  | 90  | .   | .   | 50  | 55  | .   | .  |
| <i>Lactuca orientalis</i>                               | .   | .   | .   | 73  | 40  | 50  | 18  | 30  | .  |
| <i>Echinops chloroleucus</i>                            | .   | .   | .   | 73  | 40  | 25  | 18  | 20  | .  |
| <i>Scrophularia scabiosifolia</i>                       | 60  | 20  | .   | .   | 40  | 25  | 55  | .   | .  |
| <i>Galium spurium</i>                                   | .   | 30  | 30  | .   | 40  | 75  | 45  | .   | .  |
| <i>Nepeta olgae</i>                                     | 40  | 20  | .   | 45  | .   | .   | 45  | .   | .  |
| <i>Crambe kotschyana</i>                                | .   | .   | .   | 45  | 40  | 75  | 18  | .   | .  |
| <i>Veronica intercedens</i>                             | 30  | 60  | 10  | .   | .   | .   | 36  | .   | .  |



| Syntaxonomical unit   | 1  | 2  | 3  | 4  | 5  | 6  | 7  | 8  | 9   |
|---|----|----|----|----|----|----|----|----|-----|
| <i>Verbascum songaricum</i>                                 | 60 | 50 | 20 | .  | .  | .  | .  | .  | .   |
| <i>Eremurus olgae</i>                                       | .  | .  | 70 | 9  | .  | .  | 36 | .  | .   |
| <i>Melica inaequiglumis</i>                                 | .  | .  | .  | 18 | 30 | 50 | 36 | .  | .   |
| <i>Eremurus comosus</i>                                     | .  | 70 | .  | .  | .  | .  | 36 | .  | .   |
| <i>Ziziphora pamiroalaica</i>                               | 40 | 10 | 20 | .  | .  | .  | 36 | .  | .   |
| <i>Stipa orientalis</i>                                     | .  | .  | .  | 27 | 20 | .  | .  | 60 | .   |
| <i>Cousinia ferganensis</i>                                 | .  | .  | .  | 55 | 40 | .  | .  | .  | .   |
| <i>Atraphaxis pyrifolia</i> b                               | .  | 10 | .  | 9  | .  | 25 | 36 | .  | .   |
| <i>Vicia kokanica</i>                                       | .  | .  | .  | .  | .  | .  | 82 | .  | .   |
| <i>Papaver pavoninum</i>                                    | 30 | .  | 10 | .  | .  | 75 | .  | .  | .   |
| <i>Lindelofia macrostyla</i>                                | .  | .  | .  | .  | .  | 75 | 27 | .  | .   |
| <i>Erodium oxvrrhynchum</i>                                 | .  | .  | .  | 18 | .  | .  | .  | 70 | .   |
| <i>Cousinia radians</i>                                     | .  | .  | .  | .  | 60 | 50 | .  | .  | .   |
| <i>Pimpinella puberula</i>                                  | .  | 10 | 60 | .  | .  | .  | .  | .  | .   |
| <i>Polygonum polycnemoides</i>                              | .  | 30 | 10 | .  | .  | .  | 27 | .  | .   |
| <i>Rheum maximoviczii</i>                                   | .  | .  | .  | .  | 20 | 25 | 36 | .  | .   |
| <i>Cousinia mulgediifolia</i>                               | 20 | .  | 50 | .  | .  | .  | .  | .  | .   |
| <i>Ziziphora interrupta</i>                                 | .  | .  | .  | .  | .  | .  | 45 | .  | .   |
| <i>Cephalorhynchus songaricus</i>                           | 10 | .  | .  | 9  | .  | .  | .  | .  | 22  |
| <i>Rhinopetalum bucharicum</i>                              | .  | .  | 50 | .  | .  | .  | .  | .  | .   |
| <i>Astragalus varzobicus</i>                                | .  | .  | 50 | .  | .  | .  | .  | .  | .   |
| <i>Rubia tibetica</i>                                       | .  | .  | .  | 9  | .  | 25 | 27 | .  | .   |
| <i>Linaria popovii</i>                                      | .  | .  | .  | 27 | .  | 25 | .  | .  | .   |
| <i>Pachypterygium brevipes</i>                              | .  | .  | .  | .  | .  | 50 | .  | 30 | .   |
| <i>Cousinia butkovii</i>                                    | .  | .  | .  | 36 | .  | .  | .  | .  | .   |
| <i>Colutea paulsenii</i> b                                  | .  | .  | .  | 18 | .  | .  | 18 | .  | .   |
| <i>Eremurus pubescens</i>                                   | .  | .  | 10 | .  | .  | 25 | 18 | .  | .   |
| <i>Scandix stellata</i>                                     | .  | .  | .  | .  | .  | 75 | 9  | .  | .   |
| <i>Tetrataenium olgae</i>                                   | .  | .  | .  | .  | 10 | 50 | 9  | .  | .   |
| <i>Chesneya ternata</i>                                     | .  | .  | .  | .  | .  | .  | 27 | .  | .   |
| <i>Marrubium vulgare</i>                                    | 20 | .  | .  | .  | .  | 25 | .  | .  | .   |
| <i>Glaucium fimbriigerum</i>                                | .  | 20 | 10 | .  | .  | 25 | .  | .  | .   |
| <i>Ephedra heterosperma</i> b                               | .  | .  | .  | 9  | .  | .  | .  | 20 | .   |
| <i>Stipa caucasica</i>                                      | .  | .  | .  | .  | .  | .  | .  | 10 | .   |
| <i>Glaucium squamigerum</i>                                 | .  | .  | .  | .  | .  | .  | .  | .  | 33  |
| <i>Cousinia pseudoshisakensis</i>                           | 10 | .  | .  | .  | .  | .  | .  | .  | .   |
| <i>Scutellaria adenostegia</i>                              | 30 | .  | .  | .  | .  | .  | .  | .  | .   |
| <i>Seseli lehmannianum</i>                                  | .  | .  | .  | 9  | .  | .  | .  | .  | .   |
| <i>Incarvillea olgae</i>                                    | 20 | .  | .  | .  | .  | .  | .  | .  | .   |
| <i>Geranium regelii</i>                                     | .  | 20 | .  | .  | .  | .  | .  | .  | .   |
| <i>Rochelia cardiosepala</i>                                | .  | 20 | .  | .  | .  | .  | .  | .  | .   |
| <i>Ampelopsis vicifolia</i>                                 | .  | .  | 20 | .  | .  | .  | .  | .  | .   |
| <i>Lappula drobovii</i>                                     | .  | .  | .  | 18 | .  | .  | .  | .  | .   |
| <i>Lagochilus zeravschanicus</i>                            | .  | .  | .  | 9  | .  | .  | 9  | .  | .   |
| <b>Corydalidetum kashgaricae</b>                            |    |    |    |    |    |    |    |    |     |
| <i>Corydalis kashgarica</i>                                 | .  | .  | .  | .  | .  | .  | .  | .  | 100 |
| <b>Parietarium judaicae</b>                                 |    |    |    |    |    |    |    |    |     |
| <i>Parietaria judaica</i>                                   | .  | .  | .  | .  | .  | .  | .  | 20 | 100 |
| <b>Tortulo-Cymbalarietalia &amp; Parietarietea judaicae</b> |    |    |    |    |    |    |    |    |     |
| <i>Capparis spinosa</i>                                     | .  | .  | .  | 45 | 30 | .  | .  | 50 | .   |
| <b>Asplenietea</b>  |    |    |    |    |    |    |    |    |     |
| <i>Carex koshevníkovii</i>                                  | 30 | .  | .  | .  | .  | .  | .  | .  | .   |
| <i>Artemisia rutifolia</i>                                  | .  | .  | .  | .  | .  | .  | 18 | .  | .   |
| <i>Spiraea baldshuanica</i> b                               | 20 | .  | .  | .  | .  | .  | .  | .  | .   |
| <i>Poa relaxa</i>   | 10 | .  | .  | .  | .  | .  | 9  | .  | .   |
| <i>Stipa zeravschanica</i>                                  | .  | .  | .  | .  | .  | .  | .  | 20 | .   |

ment S1). The most frequent species co-occurring within the association plots were *Bromus tectorum*, *Callipeltis cucullaris*, *Nepeta olgae*, *Piptatherum songaricum*, *Scrophularia scabiosifolia*, *Trichodema incanum*, *Vicia kokanica* and *Origanum tyttanthum*.

#### 7. *Zygophylletum atriplicoidis* ass. nova

Typus relevé: Supplement S2, rel. 12, holotypus hoc loco

Diagnostic species: *Zygophyllum atriplicoides*

The *Zygophylletum atriplicoidis* has been spotted sporadically in the central section of Jagnob and Zeravshan River valleys on southern slopes of Turkestan and Zeravshan Mts (Fig. 6, 7). The diagnostic species has an Irano-Turanian distributional type and is known from Caucasus foothills, Iran, northern Afghanistan, Tajikistan and western Kyrgyzstan (OVCHINNIKOV & KINZIKAeva 1981). The association prefers the driest sites in the colline and montane altitudinal zones at elevations of ca. 1,550 m a.s.l. (Fig. 4). The *Zygophylletum atriplicoidis* occurs on coarse-gravel and pebble deposits with very scarce soil ingredient. Plots of the association were found on open, highly radiated and warm habitats. The community inhabits the limestone and dolomite screes of alkaline reaction (pH 8.2–8.5). It was observed generally on southern expositions with very steep inclinations (mean approx. 40°). The species constituting the vegetation plots are generally scarce. The cover of the herb layer generally was between 20 and 55% with mean value of approx. 45%. The association is characterised by a low number of species in comparison to other scree vegetation in Tajikistan, having from 5 to 16 species in one relevé (mean value ca. 7). The vascular plants with the highest values of constancy and abundance are: *Capparis spinosa*, *Erodium oxyrrhynchum*, *Krashennikovia ceratoides*, *Salsola gemmascens* and *Stipa orientalis*.

#### 8. *Corydalidetum kashgaricae* ass. nova

Typus relevé: Supplement S2, rel. 42, holotypus hoc loco

Diagnostic species: *Corydalis kashgarica*

*Corydalis kashgarica* is a widely distributed scree plant across the whole Middle Asia (Pamir-Alai, Tian Shan, Kunlun Mts; OVCHINNIKOV 1975). In Tajikistan the species was found on steep conglomerate screes in the central section of Zeravshan range in Pastrudaria and Jagnob River valleys (Fig. 6). The association is distributed in the upper montane zone at elevations of approx. 1,500–2,200 m a.s.l. (mean. ca. 2,000 m; Fig. 4). It was found on conglomerate steep and compact screes of undercut colluvial cones. We found the community on limestone and dolomite debris of alkaline reaction (pH 7.8–8.6). The association inhabits generally the western expositions with almost vertical inclinations of approx. 75–85° (Fig. 4). *Corydalidetum kashgaricae* is relatively species poor with 2 to 5 species per plot (mean approx. 4). In line with species scarcity is the total herb cover with values not exceeding 50% and mean less than 40%. Among the species occurring with higher constancy only *Parietaria judaica* is worth mentioning as the co-dominating plant typical for this kind of vegetation.

Several plots have not been assigned to any of the above listed associations, nor were they included in the classification scheme. This was because we could not find a well suited species diagnosis for them as the plots were almost monospecific (e.g. *Glaucium elegans*), or in our opinion they were underdeveloped during sampling (*Chondrilla gibbistris*). Some plots may rather be related to another type of vegetation, but due to still poor

knowledge about tall forbs or xerothermophilous swards we could not be sure. It is possible that communities with contribution of *Echinops nanus* belong to xeric swards and plots with *Prangos pabularia* should be classified to tall herbs. However, these opinions are only based on visual observations and should be supported in further thorough phytosociological surveys. In the case of the community with *Zygophyllum gontscharovii* we are almost sure that this species pattern is stable and also present in other locations, however, the scarce phytosociological material and the unsuitable date of sampling during the dormancy period of some species in subtropical areas of SW Tajikistan makes considerable reservations towards the classification of that community. To facilitate the further research, we put all plots in the tables and show them in the syntaxonomic synopsis (Table 1).

## 5. Discussion

Unlike the scree plant associations of the alpine and subnival belt in Tajikistan (NOWAK et al. 2015), the communities of the colline and montane zones could be divided into two groups of calcifuge and calcicole vegetation. However, the difference in terms of floristic composition is not so evident and many species occur in both types, so it seems to be not allowed to establish separate alliances and orders as it is for European scree and rock vegetation (*Thlaspietalia rotundifolii* versus *Androsacetalia alpinae* for scree vegetation or *Potentilletalia caulescentis* versus *Androsacetalia vandellii* for rock communities). In Tajikistan alkaline and acidic substrates are inhabited by an array of common species like *Alcea nudiflora*, *Bromus tectorum*, *Centaurea squarrosa*, *Callipeltis cucullaris*, *Galium spurium*, *Hypericum scabrum* and others. Obviously there are also species distinct for both habitats like *Scutellaria intermedia* for base rich and *S. adenostegia* for acidic habitats. However, the difference seems not to be strong enough to justify the establishment of separate syntaxa. Further research is needed to define at least the suballiances. This feature seems to be typical for chasmophytic vegetation of Tajikistan dominated by the quarternary outcrops of alkaline rocks. In typical rock plant communities (*Asperulo-Poion*, *Caricion koshevnikovii*), despite some evident distinction between physico-chemical characteristic of the rock substrata, the main discrimination factors influencing the species composition were altitude, crevice size and soil amount (NOWAK A. et al. 2014a, b, c). The remarkable altitudinal amplitude in Tajikistan of more than 7,000 m causes enormous differences in microclimatic conditions and consequently in erosion type, denudation dynamic and adaptation possibilities for extreme temperatures and solar radiation. In consequence, the most important discriminating factor for species composition is the altitudinal gradient (Nowak et al. 2015). This is a fairly common pattern as we know from other studies where the altitude, bedrock type and mobility of scree material are the main factors determining species composition (VALACHOVIČ et al. 1997, PAROLLY 1998, ERMAKOV et al. 2006). That is why we propose to divide the syntaxa during our pioneer research of scree vegetation in Pamir-Alai according to their altitudinal zonation. The scree vegetation of lower mountainous elevations differs remarkably from that in alpine and subnival zones (Nowak et al. 2015). In higher elevation no acidophilous screes occur and the rock particle sizes are generally larger due to younger development stages of colluvial cones. So, apparently the species composition is also different. For *Alceion nudiflorae* we can indicate many species with considerable fidelity, occurring only or almost exclusively in communities of the colline and montane altitudinal zones. There is a range of other taxa closely related to that alliance and simultaneously diagnostic for associations e.g. *Cousinia corymbosa*, *C. refracta*, *Eremostachys tadschikistanica*, *Eremurus*

*sogdianus*, *Zygophyllum atriplicoides* and others. Against our expectations we did not find species typical for rock communities of the *Caricion koshevníkovii* in considerable constancy or abundance (NOWAK et al. 2014c). A few exceptions are *Artemisia rutifolia* and *Carex koshevníkovii*. As we supposed earlier, *Scutellaria adenostegia* should rather be excluded from the typical rock communities, because it frequently occurs in both habitats – petrophytic and scree. It is worth mentioning that in lower, colline altitudes we observed a considerable share of xerothermophilous sward species, making the difference against rock vegetation even stronger. Several taxa like *Pimpinella puberula*, *Achillea biebersteinii*, *Aphanopleura capillipes*, *Avena trichophylla*, *Gontscharovia popovii*, *Heliotropium olgae*, *Hordeum bulbosae*, *Linum corymbulosum* and *Phlomis canescens* are frequently entering the scree vegetation, especially if the denudation processes protect some soil ingredient inbetween rock particles of colluvial deposits.

The altitudinal distribution of the *Alceion nudiflorae* species is concentrated between 1,200 and 1,900 m a.s.l., while the species constituting the *Ferulo foetidissimae-Vicion kokanicae* generally inhabit the alpine zone between 1,900 and 3,600 m a.s.l. (NOWAK et al. 2015). Although we explored only a relatively narrow altitudinal zone of the 7,000 m of Tajiks vertical landscape, the particular associations reveal some differences in their elevation affinity. The highest plots were sampled for community with *Glaucium elegans*. Sites of the *Corydalidetum kashgaricae* were also relatively highly elevated. Considering the range of *Corydalis kashgarica*, which comprises the arid areas of highly elevated plateaus and mountain slopes in western China, eastern Tajikistan and southern Kyrgyzstan, as well as the remarkable distinction of the species composition and habitat conditions of that association, its distinct position in the syntaxonomical system is justified. However, it is possible that the final position of the community in vertical zonation can change. The classification into the montane belt is based on the microhabitat features of the found plots in Tajikistan and partially also on the literature data from other countries. In China the species is known from altitudes of approx. 1,200–1,800 m (Flora of China; www.efloras.org.) and from Kyrgyzstan from maximum elevation of ca. 2,100 m (PACHOMOVA 1974). So, despite the considerably high altitude of the sampled plots and the assigned range of the species in Tajikistan up to 2,400 m a.s.l. (OVCHINNIKOV 1975) we decided to include the association in the montane group. Further studies in whole Middle Asia should clarify the vertical range of *Corydalidetum kashgaricae* as well as other types of scree vegetation of the eastern, arid Pamirian geobotanical subregion. Our first insights on screes around Khorogh, Murgab and Kara-Kul Lake revealed a significant uniqueness of this area in terms of species composition of scree phytocoenoses. As examples of abundant species inhabiting the colluvial cones of eastern Pamir *Acantholimon diapensioides*, *A. velutinum*, *Artemisia fedtschenkoana*, *A. hedinii*, *A. kuscha-kewiczii*, *Astragalus alichurii*, *A. chomutovii*, *Braya pamirica*, *Hackelia murgabica*, *H. popovii* and several others might be mentioned. These taxa probably constitute considerably different, species poor associations of high and deserted areas of Tibet, Kunlun Mts and southern Tian Shan.

The lowest locations and narrow altitudinal amplitude was found for community of *Zygophyllum gontscharovii*. This is the endemic plant community of south-western Pamir-Alai, probably comprising the borderland of three states: Tajikistan, Uzbekistan and Afghanistan. Further sampling is needed to find the relation of this phytocoenosis to subtropical scree and shrub vegetation in Afghan mountains and check the final position of that community in the syntaxonomic system. The other associations are generally distributed in the typical montane

zone as far as the western Pamir-Alai mountains are concerned with only few plots ascending to the alpine belt (e.g. from *Eremostachyetum tadshikistanicae*) or descending to the colline belt (e.g. *Eremuretum sogdianae* or *Cousinietum corymbosae*).

With exclusion of the *Corydalidetum kashgaricae*, due to the species composition, the described alliance and all communities have been classified within the *Sileno brahuicae-Scutellarietalia intermediae* and *Artemisia satolinifoliae-Berberidetea sibiricae*. As discussed in NOWAK et al. (2015), the most suitable and comprehensive solution for the classification of alpine scree vegetation in Tajikistan is to include it into the Central Asian class of screes proposed by ERMAKOV et al. (2006). Examples of common species occurring in both areas are *Artemisia persica*, *A. santolinifolia*, *A. dracunculus* and *Stipa orientalis*. Other possibilities like *Thlaspietea rotundifolii* designated for Europe, *Lamio tomentosio-Chaerophylletea humilis* for Caucasus or *Heldreichietae* proposed for Taurus Mts in Turkey have to be rejected as they are remarkably distinct in species composition with almost no common species or even genera shared with Tajk's vegetation (VALACHOVIČ et al. 1997, QUEZEL 1973, PAROLLY 1998, BELONOVSKAYA 2012, BELONOVSKAYA et al. 2014). However, the southern Siberian screes, despite some microhabitat and climatic similarities (climate continentality, considerable fluctuations in temperature, aridity of the steppe and forest-steppe), are sufficiently distinct to constitute a separate order *Sileno brahuicae-Scutellarietalia intermediae*. This vegetation unit is defined by a considerable abundance of species of Irano-Turanian distributional types and consists of several *Apiaceae* genera (e.g. *Ferula* sp., *Angelica* sp., *Semenovia* sp., *Seseli* sp., *Aulacospermum* sp., *Tetrataenium olgae*, *Mediasia macrophylla* and others), *Asteraceae* (e.g. *Cousinia* sp.), *Amaryllidaceae* (*Allium* sp.), dwarf shrubs (e.g. *Artemisia* sp.) and shrubs (e.g. *Rosa* sp., *Ephedra* sp., *Cerasus* sp., *Atraphaxis* sp., *Berberis* sp.). Important components of Pamir-Alai scree vegetation also include taxa of the *Rheum*, *Euphorbia*, *Piptatherum*, *Scutellaria*, *Scrophularia* and *Zygophyllum* genera.

Mountains and rocky habitats are often centres of vascular plant endemism due to ecological separateness, high speciation rate and specific and diversified microhabitat conditions (MÉDAIL & VERLAQUE 1997, NOROOZI et al. 2008, 2011, SPEZIALE & EZCURRA 2012). This habitat and floristic uniqueness considerably influences the diversification of plant communities. This results in floristically well-defined syntaxa of regional distribution with characteristic species combinations of high fidelity (VALACHOVIČ et al. 1997, DEIL et al. 2008). As we observed in Middle Asia, the number of defined syntaxa among rock vegetation is five-fold higher than in segetal plant communities (NOWAK S. et al. 2013a, b, NOWAK S. & NOWAK A. 2013). The separateness of scree habitat is also fairly high in Tajikistan due to extreme draughtness, very harsh microclimate in higher elevation, unstable ground and frequent stone avalanches destructive to plants. Therefore, it is not surprising that on screes in Tajikistan many endemic species occur, e.g. *Allium crystallinum*, *Astragalus varzobicus*, *Cousinia butkovii*, *C. corymbosa*, *C. outichaschensis*, *C. pseudoshisakensis*, *C. refracta*, *Delphinium decoloratum*, *Eremurus olgae*, *E. comosus*, *E. pubescens*, *Ferula equisetacea*, *F. koso-polianskyi*, *Lagochilus kschtutensis*, *Rhinopetalum bucharicum* and others. This results in narrow distributional ranges of the scree plant communities of the montane and colline belt in Tajikistan. Several of them, mainly *Eremostachyetum zeravschanicae*, *Cousinietum corymbosae* and *Cousinietum refractae* are stenochoric within the country. Probably also the community with *Zygophyllum gontscharovii* is fairly restricted to the south-western part of Tajikistan or marginally reaches northern Afghanistan and south-eastern Uzbekistan.

## 6. Concluding remarks

The present survey increases the knowledge about scree vegetation of the western Pamir Alai Mts. On the basis of floristic composition and habitat features we were able to describe seven new associations, two subassociations and one alliance. Insights into screes of eastern Pamir and maybe some highly elevated sections of Vanch, Rushan, Huf, Yazgulem and Iskashim valley may further enrich the syntaxonomic system of scree vegetation in Middle Asia. These areas require further geobotanical research. The distinctiveness of the eastern Pamirian flora, especially from *Hackelia*, *Lappula*, *Nepeta*, *Acantholimon* and *Astragalus* genera, should have reflection in the diversity of scree vegetation, so the phytosociological study in this area of Tajikistan must be continued.

## Erweiterte deutsche Zusammenfassung

**Einleitung** – Tadschikistan ist ein Gebirgsland, das von alpinen Landschaften mit riesigen Schuttfächern, Schwemmkegeln, Gletschern und Moränen geprägt ist. Im vorliegenden Beitrag wird ein erster Versuch unternommen, die Vielfalt der Schuttgesellschaften in der kollinen und montanen Stufe des Pamir-Altai-Systems zu klassifizieren. Wir möchten damit folgende Fragen beantworten: (1) Welche Pflanzengesellschaften können auf diesen Standorten unterschieden werden? (2) Was sind die Standortbedingungen der gefundenen Gesellschaften? (3) Was ist ihre Artenzusammensetzung und Struktur? (4) Welche sind ihre diagnostischen Arten? Wie schon unsere früheren Studien über Felsspaltenvegetation (NOWAK A. et al. 2014a, b) stellt diese Arbeit einen Beitrag zu einem Projekt dar, an dessen Ende die syntaxonomische Klassifikation der gesamten Fels- und Schuttvegetation Mittelasiens (oder zumindest des Pamir-Alai- und Tian-Shan-Gebirges) stehen soll.

**Untersuchungsgebiet** – Die pflanzensoziologischen Aufnahmen erfolgten im westlichen Teil Tadschikistans im Pamir-Alai-Gebirge (Abb. 1 und 2). Tadschikistan liegt in Mittelasien auf einer östlichen Länge von 36°40'–41°05' und einer nördlichen Breite von 67°31'–75°14'. Es hat eine Fläche von ca. 143.500 km<sup>2</sup>. Mehr als 50 % der Landesfläche liegen oberhalb von 3.000 m Seehöhe, und mehr als 93 % liegen oberhalb von 1.000 m. Der südliche Teil Tadschikistans wird vom subtropischen Klima beeinflusst, während der nördliche Teil in der temperaten Klimazone liegt. Das Gebiet ist durch starke Sonneneinstrahlung, große jahreszeitliche Temperaturunterschiede und geringe Niederschläge gekennzeichnet.

**Methoden** – In den Jahren 2010 bis 2014 wurden insgesamt 105 Vegetationsaufnahmen in Schuttgesellschaften der kollinen und montanen Stufe des Pamir-Alai-Gebirges durchgeführt. Die Flächengröße betrug 30 m<sup>2</sup> mit Ausnahme der Bestände mit *Corydalis kashgarica*, wo aufgrund der abweichenden Physiognomie Flächen von 5 m<sup>2</sup> aufgenommen wurden. Um einen ersten Überblick über die Datenstruktur zu erhalten, wurden die Aufnahmen mit dem modifizierten TWINSPAN-Algorithmus (ROLEČEK et al. 2009) klassifiziert. Zusätzlich wurde eine DCA-Ordination durchgeführt, um die Klassifikation zu überprüfen und die Beziehungen zwischen den Einheiten herauszuarbeiten. Die Nomenklatur der Arten folgt hauptsächlich CHEREPANOV (1995) sowie Plant List (<http://www.theplantlist.org>; accessed 15 May 2015).

**Ergebnisse und Diskussion** – Es konnten acht Assoziationen unterschieden werden: *Cousinetum corymbosae*, *Eremostachyetum tadschikistanicae*, *Cousinetum refractae*, *Caccinietum dubiae*, *Eremoretum sogdiani* (mit den Subassoziationen *typicum* und *delphinietosum decolorati*), *Feruletum kuhistanicae*, *Zygophylletum atriplicoidis* und *Corydalidetum kashgaricae* (Abb. 3–6, Tab. 1, Beilage S1–2). Aufgrund ihrer floristischen Zusammensetzung fassen wir alle Gesellschaften mit Ausnahme der letztgenannten in einem neuen Verband *Alceion nudiflorae* zusammen, welchen wir zu der provisorischen Ordnung *Sileno brahuicae-Scutellarietalia intermediae* innerhalb der Klasse *Artemisio santolinifoliae-Berberidetea sibiricae* stellen. Diese Klasse wurde aus Südsibirien beschrieben (ERMA-

KOV et al. 2006) und ist durch eine Häufung von Arten mit Irano-Turanischer Verbreitung gekennzeichnet. Hierzu zählen krautige Arten der *Apiaceae*, *Asteraceae* und *Amarylidaceae* sowie Zwergsträucher (z. B. *Artemisia* sp.) und Sträucher (z. B. *Rosa* sp., *Ephedra* sp., *Cerasus* sp., *Atraphaxis* sp., *Berberis* sp.). Wichtige Komponenten der Schuttvegetation im Pamir-Alai sind außerdem Vertreter der Gattungen *Rheum*, *Euphorbia*, *Piptatherum*, *Scutellaria*, *Scrophularia* und *Zygophyllum*.

Die Artenzusammensetzung der untersuchten Gesellschaften scheint hauptsächlich durch die folgenden Faktoren gesteuert zu werden: Mobilität und Größe der Schuttpartikel, Seehöhe und Hangneigung. Im Gegensatz zur Schuttvegetation der alpinen und subnivalen Stufe Tadschikistans (vgl. NOWAK et al. 2015) können die Gesellschaften der kollinen und montanen Stufe nach dem Chemismus des Substrats in eine calcifuge und eine calcicole Gruppe unterteilt werden. Allerdings ist diese Untergliederung nicht sehr scharf, und viele Arten treten in beiden Gruppen auf, so dass es nicht möglich scheint, diese auf höherem syntaxonomischen Niveau zu trennen, analog zur europäischen Schutt- und Felspaltenvegetation (*Thlaspietalia rotundifolii* versus *Androsacetalia alpinae*, *Potentilletalia caulescentis* versus *Androsacetalia vandellii*). Das Zentrum der Verbreitung des Verbands *Alceion nudiflorae* liegt zwischen 1.200 und 1.900 m Seehöhe, während der alpine Verband *Ferulo foetidissimae-Vicion kokanicae* meist zwischen 1.900 und 3.600 m auftritt (NOWAK et al. 2015).

## Acknowledgements

We are very grateful to Prof. Adam Bodzioch for identification of rock samples. The authors wish also to thank Firuza Illarionova from the Nature Protection Team Dushanbe for assistance and help in organizing expeditions.

## Supplements

**Supplement S1.** Communities of screes of the montane and colline zones in western Pamir-Alai Mts in Tajikistan. Part 1: *Cousinietum corymbosae*, *Eremostachyetum tadschikistanicae*, *Cousinietum refractae*, *Caccinietum dubiae*, *Eremuretum sogdiani*, *Eremuretum sogdiani delphinietosum decolorati*.

**Beilage S1.** Die Gesellschaften der montanen und kollinen Schuttvegetation in Tadschikistan. Teil 1.

**Supplement S2.** Communities of screes of the montane and colline zones in western Pamir-Alai Mts in Tajikistan. Part 2: *Feruletum kuhistanicae*, *Zygophylletum atriplicoidis*, comm. with *Echinops nanus*, comm. with *Glaucium elegans*, comm. with *Prangos pabularia*, comm. with *Zygophyllum gontscharovii*, comm. with *Chondrilla gibbistrostris*, *Corydaliidetum kashgaricae*.

**Beilage S2.** Die Gesellschaften der montanen und kollinen Schuttvegetation in Tadschikistan. Teil 2.

## References

- BELONOVSKAYA, E.A. (2012): Classification of the plant communities on screes in the alpine belt of the northern Caucasus. – *Izvestia Samarskovo centra Rossijskoi Akademii Nauk* 14: 967–970.
- BELONOVSKAYA, E.A., MUCINA, L. & THEURILLAT, J.-P. (2014): Syntaxonomic and nomenclatural notes of the scree vegetation of Caucasus. – *Hacquetia* 13: 279–284.
- BRAUN-BLANQUET, J. (1964): *Pflanzensoziologie. Grundzüge der Vegetationskunde*. 3<sup>rd</sup> ed. – Springer, Wien: 865 pp.
- CHEREPANOV, S.K. (1995): *Plantae Vasculares URSS*. – Nauka, Leningrad: 516 pp.
- CHUKAVINA, A.P. (1984): *Flora Tadzchikoy SSR. 7. Zontichnye – Verbenovye (Flora of the Tajiks SSR. Vol. 7. Apiaceae – Verbenaceae)* [in Russian]. – Izdatel'stvo Nauka, Leningrad: 562 pp.
- CHYTRÝ, M. (Ed). (2009): *Vegetace České republiky. 2. Ruderální, plevelová, skalní a suťová vegetace (Vegetation of the Czech Republic 2. Ruderal, weed, rock and scree vegetation)* [in Czech, with English summaries]. – Academia, Praha: 524 pp.
- COWAN, P.J. (2007): Geographic usage of the term Middle Asia and Central Asia. – *J. Arid. Environ.* 69: 359–363.

- DEIL, U., GALÁN de MERA, A. & ORELLANA, J.A. (2008): Rock and scree plant communities in the Serra de Monchique (SW Portugal). – Feddes Repert. 119: 556–585.
- DEIL, U. & HAMMOUMI, M. (1997): Contribution à l'étude des groupements rupicoles des Bökkoya (Littoral du Rif Central, Maroc) (Studies on the rupicolous assemblages of the Bökkoya, Littoral du Rif Central, Morocco) [in French]. – Acta Bot. Malacitana 22: 131–146.
- DIMOPOULOS, P., SÝKORA, K.V., MUCINA, L. & GEORGIADIS, T. (1997): The high-rank syntaxa of the rock-cliff and scree vegetation of the mainland Greece and Crete. – Folia Geobot. Phytotax. 32: 313–334.
- DZHURAEV, A. (1970): Bioloĥo-ecolohicheskie i zhiznnyye formy rastitelnosti pervichnyh osypiey Hissarskoho hrehta (Biology, ecology and life forms of the scree flora of the Hissar range) [in Russian]. – Doklady Akademii Nauk Tadzhikskoy SSR 13: 38–50.
- DZHURAEV, A. (1972a): Rastitelnost pervichnyh osypiey zapovednika "Ramt" (Scree vegetation of the Ramit national park) [in Russian]. – Doklady Akademii Nauk Tadzhikskoy SSR 15: 85–97.
- DZHURAEV, A. (1972b): Rastitelnost pervichnyh osypiey ushc'helya Takob (Hissarskij hrebet) (Scree vegetation of the Takob valley) [in Russian]. – Doklady Akademii Nauk Tadzhikskoy SSR 15: 44–46.
- ENGLISCH, T., VALACHOVIČ, M., MUCINA, L., GRABHERR, G. & ELLMAUER, T. (1993): *Thlaspietea rotundifolii*. – In: GRABHERR, G. & MUCINA, L. (Eds.): Die Pflanzengesellschaften Österreichs. Teil II: 276–342. Fischer, Jena.
- EREN, Ö., GÖKÇEOĞLU, M. & PAROLLY G. (2004): The flora and vegetation of Bakirli Dađı (Western Taurus Mts, Turkey), including annotations on critical taxa of the Taurus range. – Willdenowia 34: 463–503.
- ERMAKOV, N., CHYTRÝ, M. & VALACHOVIČ, M. (2006): Vegetation of the rock outcrops and screes in the foreststeppe and steppe belts of the Altai and Western Sayan Mts., southern Siberia. – Phytocoenologia 36: 509–545.
- ERMOLAEVA, O.Yu. (2007): The petrophyte plant communities of high mountain limestone area, the West Caucasus. – Veg. Russ. 10: 23–37.
- GIAM, X., BRADSHAW, C.J.A., TAN, H.T.W. & SODHI, N.S. (2010): Future habitat loss and the conservation of plant biodiversity. – Biol. Conserv. 143: 1594–1602.
- GOLUB, V.B., GRECHUSHKINA, N.A., SOROKIN, A.N. & NIKOLAYCHUK, L.F. (2009): Plant communities on rock outcrops in the northwest part of the Black Sea Caucasian Coast. – Veg. Russ. 14: 3–14.
- HEIN, P., KÜRSCHNER, H. & PAROLLY, G. (1998): Phytosociological studies on high mountain plant communities of the Taurus mountains (Turkey). 2. Rock communities. – Phytocoenologia 28: 465–563.
- KINZIKAEVA, G.K. (1988): Flora Tadzhikskoi SSR. 9. Marenovye – Slozhnotsvetnye (Flora of the Tajiks SSR. Vol. 9. Moraceae – Asteraceae) [in Russian]. – Izdatelstvo Nauka, Leningrad: 568 pp.
- KOCHKAREVA, T.F. (1986): Flora Tadzhikskoy SSR. 8. Kermekovye – Podorozhnikovye (Flora of the Tajiks SSR. Vol. 8. *Limoniaceae* – *Plantaginaceae*) [in Russian]. – Izdatelstvo Nauka, Leningrad: 519 pp.
- KOSIŃSKI, M. (2001): Synecological differentiation of the scree vegetation of the Polish Tatra Mountains. – Phytocoenologia 31: 499–520.
- KOVALEVSKAYA, S.S. (1971): Conspectus florum Asiae Mediae. Vol. 2 (Conspect of the flora of Middle Asia. Vol. 2) [in Russian]. – Izdatelstvo FAN Uzbekskoi SSR. Tashkent: 360 pp.
- LATIPOVA, W.A. (1968): Kolichestvo osadkov. – In: NARZIKULOV, I.K. & STANJUKOVICH, K.W. (Eds.): Atlas Tajikskoi SSR (Precipitation, in Atlas of the Tajiks SSR) [in Russian]: 68–69. Akademia Nauk Tajikskoi SSR, Dushanbe-Moskva.
- MÉDAIL, F. & VERLAQUE, R. (1997): Ecological characteristics and rarity of endemic plants from southeastern France and Corsica: implications for biodiversity conservation. – Biol. Conserv. 80: 269–281.
- MITTERMEIER, R.A., GIL, P.R., HOFFMAN, M., PILGRIM, J., BROOKS, T., GOETTSCHE-MITTERMEIER, C., LAMOREUX, J. & Da FONSECA, G.A.B. (2005): Hotspots revisited: Earth's biologically richest and most threatened terrestrial ecoregions. – Conservation International and Agrupacion Sierra Madre, Monterrey, Mexico: 392 pp.
- MOTA POVEDA, J.F., GÓMEZ MERCADO, F. & VALLE TENDERO, F. (1991): Rupicolous vegetation of the betic ranges (south Spain). – Vegetatio 94: 101–113.



- MUCINA, L., VALACHOVIČ, M., JAROLÍMEK, I., ŠEFFER, J., KUBINSKÁ, A. & PIŠŮT, I. (1990): The vegetation on rock fissures, screes, and snow-beds in the Pirin planina Mountains (Bulgaria). – Stud. Geobot. 10: 15–58.
- NARZIKULOV, I.K. & STANJUKOVICH, K.W. (1968): Atlas Tajikskoi SSR.(Atlas of the Tajiks SSR) [in Russian]. – Akademia Nauk Tajikskoi SSR, Dushanbe-Moskva.
- NEDZVEDSKIY, A.P. (1968): Geologicheskoe stroenye. – In: NARZIKULOV, I.K., STANJUKOVICH, K.W. (Eds.): Atlas Tajikskoi SSR (Geology, in Atlas of the Tajiks SSR) [in Russian]: 14–15. Akademia Nauk Tajikskoi SSR, Dushanbe-Moskva.
- NOBIS, M. (2011a): Remarks on the taxonomy and nomenclature of the *Stipa tianshanica* complex (Poaceae), on the base of a new record for the flora of Tajikistan (central Asia). – Nord. J. Bot. 29: 194–199.
- NOBIS, M. (2011b): *Stipa ×brozhiana* (Poaceae) nothosp. nov. from the western Pamir Alai Mts (middle Asia) and taxonomical notes on *Stipa ×tzvelevii*. – Nord. J. Bot. 29: 458–464.
- NOBIS, M. (2013): Taxonomic revision of the *Stipa lipskyi* group (Poaceae: *Stipa* section *Smirnovia*) in the Pamir Alai and Tian-Shan Mountains. – Plant Syst. Evol. 299: 1307–1354.
- NOBIS, M., EBEL, A.L., NOWAK, A., TURGINOV, O.T., KUPRIYANOV, A.N., NOBIS, A., OLONOVA, M.V., PASZKO, B., PIWOWARCZYK, R., CHEN, W-L, GUDKOVA, P.D., KLICHOWSKA, E., NOWAK, S. & PUJADAS-SALVÀ, A.J. (2014b): Contribution to the flora of Asian and European countries: new national and regional vascular plant records. 2. – Acta Bot. Gallica 161: 209–221.
- NOBIS, M., NOBIS, A., NOWAK, A. & NOWAK, S. (2014a): A new variety of *Diarthron* (*Thymelaeaceae*) from Tajikistan. – Plant Biosyst. 148: 1032–1034.
- NOBIS, M. & NOWAK, A. (2011): New data to the vascular flora of the central Pamir Alai Mountains (Tajikistan, middle Asia). – Pol. Bot. J. 56: 195–201.
- NOBIS, M., NOWAK, A., EBEL, A.L., NOBIS, A., NOWAK, S., GUDKOVA, P.D., VERKHOZINA, A.V., ERST, A.S., LAZARSKI, G., OLONOVA, M.V., PIWOWARCZYK, R., BOBROV, A.A., KHRUSTALEVA, I.A., PLÁŠEK, V., SILANTYEVA, M.M. & ZALEWSKA-GALOSZ, J. (2015): New National and Regional Vascular Plants Records 3. – Acta Bot. Gallica 162: 103–115.
- NOBIS, M., NOWAK, A. & NOBIS, A. (2013): *Stipa zeravshanica* (Poaceae) an endemic species from rocky walls of the western Pamir Alai Mountains (middle Asia). – Nord. J. Bot. 31: 666–675.
- NOROOZI, J., AKHANI, H. & BRECKLE, S.-W. (2008): Biodiversity and phytogeography of the alpine flora of Iran. – Biodivers. Conserv. 17: 493–521.
- NOROOZI, J., PAULI, H., GRABHERR, G. & BRECKLE, S.-W. (2011): The subnival–nival vascular plant species of Iran: a unique high-mountain flora and its threat from climate warming. – Biodivers. Conserv. 20: 1319–1338.
- NOROOZI, J., WILLNER, W., PAULI, H. & GRABHERR, G. (2014): Phytosociology and ecology of the high-alpine to subnival scree vegetation of N and NW Iran (Alborz and Azebaijan Mts.). – Appl. Veg. Sci. 17: 142–161.
- NOWAK, A. & NOBIS, M. (2012): Distribution patterns, floristic structure and habitat requirements of the alpine river plant community *Stuckeniaetum amblyphyllae* ass. nova (*Potametea*) in the Pamir Alai Mountains (Tajikistan). – Acta Soc. Bot. Pol. 80: 235–238.
- NOWAK, A. & NOBIS, M. (2013): Distribution, floristic structure and habitat requirements of the riparian forest community *Populeetum talassicae* ass. nova in the Central Pamir-Alai Mts (Tajikistan, Middle Asia). – Acta Soc. Bot. Pol. 82: 47–55.
- NOWAK, A., NOBIS, M., NOWAK, S. & NOBIS, A. (2014c): Distribution, ecology and conservation status of *Dionysia involucrata* Zapr., an endangered endemic of Hissar Mts (Tajikistan, Middle Asia). – Acta Soc. Bot. Pol. 83: 123–135.
- NOWAK, A., NOWAK, S. & NOBIS, M. (2011): Distribution patterns, ecological characteristic and conservation status of endemic plants of Tadzhikistan – A global hotspot of diversity. – J. Nat. Conserv. 19: 296–305.
- NOWAK, A., NOWAK, S. & NOBIS, M. (2013): Weed communities of rice fields in the central Pamir Alai Mountains (Tajikistan, Middle Asia). – Phytocoenologia 43: 101–126.
- NOWAK, A., NOWAK, S. & NOBIS, M. (2014b): Vegetation of soil rock faces and fissures of the alpine and subnival zone in the Pamir Alai Mountains (Tajikistan, Middle Asia). – Phytocoenologia 44: 81–104.
- NOWAK, A., NOWAK, S., NOBIS, M. & NOBIS, A. (2014a): Vegetation of rock clefts and ledges in the Pamir Alai Mts, Tajikistan (Middle Asia). – Cent. Eur. J. Biol. 9: 444–460.

- NOWAK, A., NOWAK, S., NOBIS, M. & NOBIS, A. (2014d): Vegetation of rock crevices of the montane and colline zones in the Pamir-Alai and Tian Shan Mts in Tajikistan (Middle Asia). – *Plant Biosyst.* 148: 1199–1210.
- NOWAK, A., NOWAK, S., NOBIS, M. & NOBIS, A. (2015): Vegetation of taluses and screes of the alpine zone in the Pamir Alai Mountains (Tajikistan, Middle Asia). – *Phytocoenologia* 45: 299–324.
- NOWAK, S. & NOWAK, A. (2013): Weed communities of root crops in the Pamir Alai Mts, Tajikistan (Middle Asia). – *Acta Soc. Bot. Pol.* 82: 135–146.
- NOWAK, S., NOWAK, A. & NOBIS, M. (2013a): Weed communities of rice fields in the central Pamir Alai Mountains (Tajikistan, Middle Asia). – *Phytocoenologia* 43: 101–126.
- NOWAK, S., NOWAK, A., NOBIS, M. & NOBIS, A. (2013b): Weed vegetation of cereal crops in Tajikistan (Pamir Alai Mts, Middle Asia). – *Phytocoenologia* 43: 225–243.
- ORTIZ, S. & RODRIGUEZ-OUBINA, J. (1993): Synopsis of the rupicolous vegetation of Galicia (North-western Iberian Peninsula). – *Folia Geobot. Phytotax.* 28: 15–49.
- OVCHINNIKOV, P.N. (1957): Flora Tajikskoi SSR. 1, Paprotnikoobraznye – Zlaki (Flora of the Tajiks SSR. Vol. 1. *Pteridophyta – Poaceae*) [in Russian]. – Izdatelstvo Akademii Nauk SSSR, Moskva – Leningrad: 544 pp.
- OVCHINNIKOV, P.N. (1963): Flora Tadzhikskoy SSR. 2, Osokovye – Orkhidnye (Flora of the Tajiks SSR. Vol. 2. *Cyperaceae – Orchidaceae*) [in Russian]. – Izdatelstvo Akademii Nauk SSSR, Moskva – Leningrad: 456 pp.
- OVCHINNIKOV, P.N. & KINZIKAEVA, G.K. (1981): Flora Tajikskoi SSR. Vol. 6, Bobovye (rod Astragal) – Tsinomorievye (Flora of the Tajiks SSR. Vol. 6. *Fabaceae – genus Astragalus*) [in Russian]. – Izdatelstvo Nauka, Leningrad: 725 pp.
- PACHOMOVA, M.G. (1974): Conspectus florum Asiae Mediae. Vol. 4 (Conspect of the flora of Middle Asia. Vol. 4) [in Russian]. – Izdatelstvo FAN Uzbekskoi SSR, Tashkent: 269 pp.
- PAROLLY, G. (1998): Phytosociological studies on high mountain plant communities of the South Anatolian Taurus mountains. 1. Scree plant communities (*Heldreichietea*): A synopsis. – *Phytocoenologia* 28: 233–284.
- PIGNATTI, S., OBERDORFER, E., SCHAMINÉE, J.H.J. & WESTHOFF, V. (1995): On the concept of vegetation class in phytosociology. – *J. Veg. Sci.* 6: 143–152.
- QUÉZEL, P. (1973): Contributions à l'étude phytosociologiques du massif du Taurus. – *Phytocoenologia* 1: 131–222.
- RASULOVA, M.R. (1991): Flora Tadzhikskoy SSR. 10, Slozhnotsvetnye (Flora of the Tajiks SSR. Vol. 10. *Asteraceae*) [in Russian]. – Izdatelstvo Nauka, Leningrad: 617 pp.
- ROLEČEK, J., TICHÝ, L., ZELENÝ, D. & CHYTRÝ, M. (2009): Modified TWINSpan classification in which the hierarchy respects cluster heterogeneity. – *J. Veg. Sci.* 20: 596–602.
- SAFAROV, N. (2003): National strategy and action plan on conservation and sustainable use of biodiversity. – Governmental Working Group of the Republic of Tajikistan. Dushanbe: 199 pp.
- SANDA, V., ÖLLERER, K. & BURESCU, P. (2008): Fitocenozele din România (Phytocoenoses of Romania) [in Romanian]. – *Ars Docendi, Universitatea din Bucuresti, Bucuresti*: 570 pp.
- SPEZIALE, K.L. & EZCURRA, C. (2012): The role of outcrops in the diversity of Patagonian vegetation: Relicts of glacial palaeofloras? – *Flora* 207: 141–149.
- TER BRAAK, C.J.F. & ŠMILAUER, P. (2002): CANOCO reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). – Microcomputer Power, Ithaca, New York: 500 pp.
- TICHÝ, L. (2002): JUICE, software for vegetation classification. – *J. Veg. Sci.* 13: 451–453.
- VALACHOVIČ, M., DIERSSEN, K., DIMOPOULOS, P., HADAČ, E., LOIDI, J., MUCINA, L., ROSSI, G., VALLE TENDERO, F. & TOMASELLI, M. (1997): The vegetation of screes – A synopsis of higher syntaxa in Europe. – *Folia Geobot. Phytotax.* 32: 173–192.
- VLADIMIROVA, W.N. (1968): Tipy klimatov. – In: NARZIKULOV, I.K. & STANJUKOVICH, K.W. (Eds.): Atlas Tajikskoi SSR. (Climatic types, Atlas of the Tajiks SSR) [in Russian]: 33–35. Akademia Nauk Tajikskoi SSR, Dushanbe-Moskva.
- WEBER, H.E., MORAVEC, J. & THEURILLAT, J.P. (2000): International code of phytosociological nomenclature. 3<sup>rd</sup> ed. – *J. Veg. Sci.* 11: 739–768.
- WESTHOFF, V. & VAN DER MAAREL, E. (1973): The Braun-Blanquet approach. – In: WHITTAKER, R.H. (Ed.): Handbook of Vegetation Science 5, Classification and ordination of communities: 619–726. Junk, The Hague.





# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Tuexenia - Mitteilungen der Floristisch-soziologischen Arbeitsgemeinschaft](#)

Jahr/Year: 2016

Band/Volume: [NS\\_36](#)

Autor(en)/Author(s): Nowak Arkadiusz, Nowak Sylwia, Nobis Marcin, Nobis Agnieszka

Artikel/Article: [Vegetation of screes of the montane and colline zones in the Pamir-Alai Mts in Tajikistan \(Middle Asia\) 223-248](#)