## E-Z-Flo Dichlone 3

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

DIRECTIONS FOR USE

635-191

**ACTIVE INGREDIENTS:** 

Dichlone (2,3-Dichloro-1,4-Naphthoquinone)

3.00%

**INERT INGREDIENTS:** 

97.00%

Total

100.00%

DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Net Contents and Wahufacturer
On Container

**WARNING** 

Dichlone is a skin irritant, especially during warm weather. Avoid prolonged or repeated skin contact especially during warm weather. Irritation is similar to sunburn, evidence fails to show any permanent injury. Avoid inhalation of dust. Wear protective clothing. Wash hands and face thoroughly with soap and water after use, before smoking or eating. Keep away from feed or food products. Protective creams are suggested for the hands and faces of sensitive individuals when dusting with Dichlone.

Keep out of any body of water. Fish may be killed.

Use only as directed: Apply Dichlone 3 Dust at the rate of 30-40 pounds per acre to the crops listed below unless otherwise indicated.

## RECOMMENDATIONS

APPLES—For control of apple scab, black rot and cedar apple rust from delayed dormant through 10 days after petal fall—Begin applications at the first sign of green tissue on developing buds and repeat applications during rains or immediately after as conditions warrant through 10 days after petal fall. Due to combined protective and eradicative action, it should not be necessary to dust oftener than once every 5 days. Do not apply to apples later than one day of harvest.

PEACHES, PRUNES AND PLUMS—For control of brown rot blossom blight, apply at pink, full bloom and preshuck fall. When cool, wet weather occurs, resulting in a prolonged bloom period, apply Dichlone Dust every 4 days. Do not apply to peaches later than 7 days of harvest. Do not apply to plums and prunes later than 3 days of harvest.

SWEET AND RED TART CHERRIES—For control of brown rot blossom blight—Apply at early bloom, full bloom and petal fall. When cool, wet weather occurs resulting in a prolonged bloom period, apply every 4 days—For control of cherry leaf spot, apply at petal fall and twice more at 10 day intervals. Do not apply to cherries later than 3 days of harvest.

BEANS—Anthracnose, begin applications just before bloom and repeat at 7-10 day intervals as necessary. Do not exceed 37 lbs. per acre application. Do not apply within 7 days of harvest.

CELERY: Celery blight—Begin applications about 7 to 10 days after plants are set. Use 20-25 lbs. per acre and repeat at 7 to 10 day intervals as needed.

TOMATOES—Late blight and gray mold, apply 15-20 lbs. per acre beginning when plants are 3 to 4 inches high and repeat as necessary